

## The Dual Rings of an R-Coring Revisited

Laurent Poinsoot & Hans-E. Porst

To cite this article: Laurent Poinsoot & Hans-E. Porst (2016) The Dual Rings of an R-Coring Revisited, Communications in Algebra, 44:3, 944-964, DOI: [10.1080/00927872.2014.990031](https://doi.org/10.1080/00927872.2014.990031)

To link to this article: <http://dx.doi.org/10.1080/00927872.2014.990031>



Published online: 25 Jan 2016.



Submit your article to this journal [↗](#)



View related articles [↗](#)



View Crossmark data [↗](#)

## THE DUAL RINGS OF AN $R$ -CORING REVISITED

Laurent Poinso<sup>1</sup> and Hans-E. Porst<sup>2</sup>

<sup>1</sup>LIPN, CNRS (UMR 7030), Université Paris 13, Sorbonne Paris Cité, Villetaneuse, France

<sup>2</sup>Department of Mathematical Sciences, University of Stellenbosch, Stellenbosch, South Africa

*It is shown that for every monoidal bi-closed category  $\mathbb{C}$  left and right dualization by means of the unit object not only defines a pair of adjoint functors, but that these functors are monoidal as functors from  $\mathbb{C}^{\text{op}}$ , the dual monoidal category of  $\mathbb{C}$  into the transposed monoidal category  $\mathbb{C}^!$ . We thus generalize the case of a symmetric monoidal category, where this kind of dualization is a special instance of convolution. We apply this construction to the monoidal category of bimodules over a not necessarily commutative ring  $R$  and so obtain various contravariant dual ring functors defined on the category of  $R$ -corings. It becomes evident that previous, hitherto apparently unrelated, constructions of this kind are all special instances of our construction and, hence, coincide. Finally, we show that Sweedler's Dual Coring Theorem is a simple consequence of our approach and that these dual ring constructions are compatible with the processes of (co)freely adjoining (co)units.*

**Key Words:** Dualization; Monoidal bi-closed categories;  $R$ -rings and  $-$ corings.

**2010 Mathematics Subject Classification:** Primary: 16T15; Secondary: 18D10.

### INTRODUCTION

The purpose of this analysis of dual rings of  $R$ -corings is twofold. First, we compare and explain conceptually the known constructions of such rings as given in [14] and [15] (see also [6]). Second, we apply the methods of [11] and prove that these dualizations are compatible with the operations of adjoining units and counits to  $R$ -rings and  $R$ -corings, respectively.

Concerning the first mentioned topic, recall that Sweedler in [14] constructs, for a given  $R$ -coring  $\mathbb{C} = (C, \Delta, \epsilon)$ , i.e., a comonoid in the monoidal category  $R^{\text{Mod}}_R$

Received October 7, 2014; Revised November 3, 2014. Communicated by A. Facchini.

To Gérard H. E. Duchamp, on the occasion of his retirement.

Address correspondence to Prof. Laurent Poinso, LIPN, CNRS (UMR 7030), Université Paris 13, Sorbonne Paris Cité, 99 av. J. B. Clément, 93430 Villetaneuse, France; E-mail: laurent.poinso@lipn.univ-paris13.fr

Second address: CReA, French Air Force Academy, Base aérienne 701, 13661 Salon-de-Provence, France.

Permanent address: Department of Mathematics, University of Bremen, 28359 Bremen, Germany.

of  $R$ - $R$ -bimodules (with tensor product  $- \otimes_R -$  over  $R$  and  $\lambda_C : R \otimes_R C \rightarrow C$  and  $\rho_C : C \otimes_R R \rightarrow C$  the canonical isomorphisms), left and right dual unital rings:

$\mathbf{Sw}_l(\mathbf{C}) := (*C, m_l, u_l)$ , with  $*C := {}_R \text{hom}(C, R)$ , the abelian group of left  $R$ -module homomorphisms from  $C$  to  $R$ , multiplication  $m_l$  acting on pairs  $(\mu, \nu) \in *C \times *C$  as  $m_l(\mu, \nu) = \mu \circ \rho_C \circ (C \otimes_R \nu) \circ \Delta$  and unital element  $u_l = \epsilon$ .

$\mathbf{Sw}_r(\mathbf{C}) := (C^*, m_r, u_r)$  with  $C^* := \text{hom}_R(C, R)$ , the abelian group of right  $R$ -module homomorphisms from  $C$  to  $R$ , multiplication  $m_r$  acting on pairs  $(\mu, \nu) \in C^* \times C^*$  as  $m_r(\mu, \nu) = \nu \circ \lambda_C \circ (\mu \otimes_R C) \circ \Delta$  and unital element  $u_r = \epsilon$ .

These rings are shown to be equipped with ring antihomomorphisms<sup>1</sup>  $\lambda_C : R \rightarrow \mathbf{Sw}_l(\mathbf{C})$  and  $\rho_C : R \rightarrow \mathbf{Sw}_r(\mathbf{C})$ , respectively, where  $\lambda_C(b)(c) = \epsilon(cb)$  and  $\rho_C(b)(c) = \epsilon(bc)$  for each  $c \in C$  and  $b \in R$ .

Thus,  $(\mathbf{Sw}_l(\mathbf{C}), \lambda_C)$  and  $(\mathbf{Sw}_r(\mathbf{C}), \rho_C)$  are in fact objects in the comma category  $R^{\text{op}} \downarrow_1 \mathbf{Ring}$ . The assignments  $\mathbf{C} \mapsto (\mathbf{Sw}_l(\mathbf{C}), \lambda_C)$  and  $\mathbf{C} \mapsto (\mathbf{Sw}_r(\mathbf{C}), \rho_C)$  are shown moreover to define (contravariant) functors from  ${}_\epsilon \mathbf{Coring} := \mathbf{Comon}(R^{\text{Mod}}_R)$  into  $R^{\text{op}} \downarrow_1 \mathbf{Ring}$ .

Recall also that Takeuchi in [15] in a more categorical way defines a contravariant functor  $\mathbf{D}_l$  from  ${}_\epsilon \mathbf{Coring}$  into the category  $\mathbf{Mon}(R^{\text{op}} \text{Mod}_R^{\text{op}})$  of monoids in the monoidal category  $R^{\text{op}} \text{Mod}_R^{\text{op}}$  of  $R^{\text{op}}$ - $R^{\text{op}}$ -bimodules (with tensor product  $- \otimes_{R^{\text{op}}} -$  over  $R^{\text{op}}$ ), where the underlying abelian group of  $\mathbf{D}_l(\mathbf{C})$  again is  $*C$ . Neither Takeuchi nor the more recent survey [6] relate the latter construction to Sweedler's.

As we are going to show in Section 2.2, both constructions are essentially the same and imply Sweedler's Dual Coring Theorem in a simple way. Here we will show, in addition (see Remark 15 (3)), how to understand conceptually a further dual ring construction given in [14] as well.

All of this is based on the purely categorical result that, given a not necessarily symmetric monoidal bi-closed category  $\mathbf{C} = (\mathbf{C}, - \otimes -, I)$  as, e.g., the category  $R^{\text{Mod}}_R$  of  $R$ - $R$ -bimodules with its standard monoidal structure, the (contravariant) left and right internal hom-functors  $[-, I]_l$  and  $[-, I]_r$  can be seen as monoidal functors (see Theorem 6). To the best of our knowledge this fact, certainly known in case  $\mathbf{C}$  is symmetric monoidal, is not known yet in the nonsymmetric case. This shows, in particular, that it is misleading to some extent to consider dualization (of coalgebras) simply as a special instance of the convolution construction: Dualization is a construction in its own right, which coincides with (a special instance) of convolution in the symmetric case.

In order to make our analysis accessible to readers not too familiar with the theory of monoidal categories, we recall its basic elements as far as they are needed.

<sup>1</sup>We here use Sweedler's original notation for these maps: Thus,  $\lambda_C$  and  $\rho_C$  here should not be mistaken for the left and right unit constraints  $\lambda_C$  and  $\rho_C$  in a monoidal category, i.e., the canonical isomorphisms mentioned above.

1. BASICS

1.1. Monoidal Categories and Functors

Throughout,  $\mathbf{C} = (\mathbf{C}, - \otimes -, I, \alpha, \lambda, \varrho)$  denotes a monoidal category with  $\alpha$  the associativity and  $\lambda$  and  $\varrho$  the left and right unit constraints. If  $\mathbf{C}$  is even symmetric monoidal, the symmetry will be denoted by  $\tau = (C \otimes D \xrightarrow{\tau_{CD}} D \otimes C)_{C,D}$ .

Recall that  $\mathbf{C}$  is called *monoidal left closed*, provided that, for each  $\mathbf{C}$ -object  $C$  the functor  $C \otimes -$  has a right adjoint  $[C, -]_l$ . If each functor  $- \otimes C$  has a right adjoint, denoted by  $[C, -]_r$ ,  $\mathbf{C}$  is called *monoidal right closed*.  $\mathbf{C}$  is called *monoidal bi-closed*, provided that  $\mathbf{C}$  is monoidal left and right closed.

The counits  $C \otimes [C, X]_l \rightarrow X$  and  $[C, X]_r \otimes C \rightarrow X$  of these adjunctions will be denoted by  $ev^l$  and  $ev^r$ , respectively. By parametrized adjunctions (see [10]) one thus has functors  $[-, -]_r$  and  $[-, -]_l : \mathbf{C}^{op} \times \mathbf{C} \rightarrow \mathbf{C}$ . In particular, for each  $X$  in  $\mathbf{C}$ , there are the contravariant functors  $[-, X]_r$  and  $[-, X]_l$  on  $\mathbf{C}$ .

For  $C \xrightarrow{f} D$  in  $\mathbf{C}$ , the morphism  $[D, X]_l \xrightarrow{[f, X]_l} [C, X]_l$  is the unique morphism such that the following diagram commutes:

$$\begin{array}{ccc} C \otimes [D, X]_l & \xrightarrow{C \otimes [f, X]_l} & C \otimes [C, X]_l \\ f \otimes [D, X]_l \downarrow & & \downarrow ev_l \\ D \otimes [D, X]_l & \xrightarrow{ev_l} & X \end{array}$$

Similarly for  $[-, X]_r$ .

In each such category there are the following situations (see, e.g., [9, Eqs. 1.25, 1.26, 1.27]):

1. Natural isomorphisms  $\text{hom}_{\mathbf{C}}(I, [C, D]_l) \simeq \text{hom}_{\mathbf{C}}(C, D) \simeq \text{hom}_{\mathbf{C}}(I, [C, D]_r)$ ;
2. Natural isomorphisms  $C \xrightarrow{j_C} [I, C]_l$  and  $C \xrightarrow{i_C} [I, C]_r$ , corresponding by adjunction to  $I \otimes C \xrightarrow{j_C} C$  and  $C \otimes I \xrightarrow{i_C} C$ , respectively;
3. Natural isomorphisms

$$[C, [D, A]_r]_r \rightarrow [C \otimes D, A]_r \quad \text{and} \quad [C, [D, A]_l]_l \rightarrow [C \otimes D, A]_l,$$

whose images under  $\text{hom}_{\mathbf{C}}(I, -)$  are, respectively, the isomorphisms

$$\begin{aligned} \text{hom}_{\mathbf{C}}(C, [D, A]_r) &\simeq \text{hom}_{\mathbf{C}}(C \otimes D, A) \quad \text{and} \\ \text{hom}_{\mathbf{C}}(D, [C, A]_l) &\simeq \text{hom}_{\mathbf{C}}(C \otimes D, A) \end{aligned}$$

expressing the adjunctions for right and left tensoring. These isomorphisms will be noted by  $\Pi_{C,D}^r$  and  $\Pi_{C,D}^l$ , respectively, in the special instance  $A = I$ ;

4. Natural transformations

$$\Theta_{C,D}^r : D \otimes [C, I]_r \rightarrow [C, D]_r \quad \text{and} \quad \Theta_{C,D}^l : [C, I]_l \otimes D \rightarrow [C, D]_l$$

corresponding by adjunction to

$$(D \otimes [C, I]_r) \otimes C \simeq D \otimes ([C, I]_r \otimes C) \xrightarrow{D \otimes \text{ev}_{C,I}^r} D \otimes I \xrightarrow{\rho_D} D$$

and, respectively,

$$C \otimes ([C, I]_l \otimes D) \simeq (C \otimes [C, I]_l) \otimes D \xrightarrow{\text{ev}_{C,I}^l \otimes D} I \otimes D \xrightarrow{\lambda_D} D.$$

Given a monoidal category  $\mathbb{C} = (\mathbb{C}, - \otimes -, I, \alpha, \lambda, \varrho)$ , there are the following simple ways of constructing new monoidal categories:

- $\mathbb{C}' = (\mathbb{C}, - \otimes' -, I, \alpha', \lambda', \rho')$  is a monoidal category with  $C \otimes' D = D \otimes C$ ,  $\alpha'_{ABC} = \alpha_{CBA}$ ,  $\rho'_C = \lambda_C$  and  $\lambda'_C = \rho_C$ .  $\mathbb{C}'$  is called the *transpose* of  $\mathbb{C}$ .
- $\mathbb{C}^{\text{op}} = (\mathbb{C}^{\text{op}}, - \otimes -, I, \alpha^{\text{op}}, \lambda^{\text{op}}, \rho^{\text{op}})$  is a monoidal category with constraints being the inverses of those in  $\mathbb{C}$ .  $\mathbb{C}^{\text{op}}$  is called the *dual* of  $\mathbb{C}$ .

By  $\mathbf{Mon}\mathbb{C}$  and  $\mathbf{Comon}\mathbb{C}$ , we denote the categories of monoids  $(M, m, e)$  in  $\mathbb{C}$  and of comonoids  $(C, \Delta, \epsilon)$  in  $\mathbb{C}$ , respectively. Omitting the units  $e$  and counits  $\epsilon$ , respectively, we obtain the categories  $\mathbf{Sgr}\mathbb{C}$  of semigroups and  $\mathbf{Cosgr}\mathbb{C}$  of co-semigroups in  $\mathbb{C}$ .

**Fact 1.** Concerning these constructions, the following facts are obvious:

1. If  $\mathbb{C}$  is symmetric monoidal, then  $\mathbb{C}$  and  $\mathbb{C}'$  are monoidally equivalent (even isomorphic).
2.  $(\mathbb{C}')^{\text{op}} = (\mathbb{C}^{\text{op}})'$ .
3.  $\mathbf{Mon}(\mathbb{C}') = \mathbf{Mon}\mathbb{C}$  and  $\mathbf{Comon}(\mathbb{C}') = \mathbf{Comon}\mathbb{C}$ .
4.  $\mathbf{Mon}\mathbb{C}^{\text{op}} = (\mathbf{Comon}\mathbb{C})^{\text{op}}$ .
5. If  $\mathbb{C}$  is monoidal bi-closed, then so is  $\mathbb{C}'$ ; its internal hom-functors can be chosen as  $[C, -]'_l = [C, -]_r$  and  $[C, -]'_r = [C, -]_l$ .

We briefly recall the following definitions and facts which are fundamental for this note.

**Definition 2.** Let  $\mathbb{C} = (\mathbb{C}, - \otimes -, I)$  and  $\mathbb{C}' = (\mathbb{C}', - \otimes' -, I')$  be monoidal categories. A *monoidal functor from  $\mathbb{C}$  to  $\mathbb{C}'$*  is a triple  $(F, \Phi, \phi)$ , where  $F : \mathbb{C} \rightarrow \mathbb{C}'$  is a functor,  $\Phi_{C_1, C_2} : FC_1 \otimes' FC_2 \rightarrow F(C_1 \otimes C_2)$  is a natural transformation, and  $\phi : I' \rightarrow FI$  is a  $\mathbb{C}$ -morphism, subject to certain coherence conditions (see, e.g., [13]). A monoidal functor is called *strong monoidal*, if  $\Phi$  and  $\phi$  are isomorphisms and *strict monoidal*, if  $\Phi$  and  $\phi$  are identities.

An *opmonoidal functor* from  $\mathbb{C}$  to  $\mathbb{C}'$  is a monoidal functor from  $\mathbb{C}^{\text{op}}$  to  $\mathbb{C}'^{\text{op}}$ .

Given monoidal functors  $\mathbb{F} = (F, \Psi, \psi)$  and  $\mathbb{G} = (G, \Phi, \phi)$  from  $\mathbb{C}$  to  $\mathbb{D}$ , a natural transformation  $\mu : F \Rightarrow G$  is called a *monoidal transformation*  $\mathbb{F} \Rightarrow \mathbb{G}$ , if the

following diagrams commute for all  $C, D$  in  $\mathbf{C}$ :

$$\begin{array}{ccc}
 FC \otimes FD & \xrightarrow{\mu_C \otimes \mu_D} & GC \otimes GD \\
 \Psi_{C,D} \downarrow & & \downarrow \Phi_{C,D} \\
 F(C \otimes D) & \xrightarrow{\mu_{C \otimes D}} & G(C \otimes D)
 \end{array}
 \qquad
 \begin{array}{ccc}
 I & \xrightarrow{\psi} & FI \\
 \searrow \phi & & \downarrow \mu_I \\
 & & GI
 \end{array}$$

A *monoidal isomorphism* is a natural isomorphism which is a monoidal transformation.

A *monoidal equivalence* of monoidal categories is given by monoidal functors  $\mathbb{F} = (F, \Psi, \psi)$  and  $\mathbb{G} = (G, \Phi, \phi)$  and monoidal isomorphisms  $\eta : \text{Id} \Rightarrow GF$  and  $\epsilon : FG \Rightarrow \text{Id}$ .

**Remarks 3** ([2]).

1. The composition of monoidal functors is a monoidal functor.
2. By a *monoidal subcategory* of a monoidal category  $\mathbf{C} = (\mathbf{C}, - \otimes -, I)$  we mean a full subcategory  $\mathbf{A}$  of  $\mathbf{C}$ , closed under tensor products and containing  $I$ . The embedding  $E$  of  $\mathbf{A}$  into  $\mathbf{C}$  then is a monoidal functor with the monoidal structure given by identities.

Let  $\mathbb{F} = (F, \Phi, \phi) : \mathbf{C} \rightarrow \mathbf{D}$  be a monoidal functor, and let  $\mathbf{C}'$  and  $\mathbf{D}'$  be monoidal subcategories of  $\mathbf{C}$  and  $\mathbf{D}$ , respectively. By a *restriction* of  $\mathbb{F}$  to these subcategories is meant a monoidal functor  $\mathbb{F}' = (F', \Phi', \phi') : \mathbf{C}' \rightarrow \mathbf{D}'$  satisfying (a)  $F'C = FC$  for all  $C$  in  $\mathbf{C}'$ , (b)  $\Phi'_{A,B} = \Phi_{A,B}$  for all  $A, B$  in  $\mathbf{C}'$ , and (c)  $\phi' = \phi$ . This is equivalent to saying that  $\mathbb{F}'\mathbb{E}_{\mathbf{C}'} = \mathbb{E}_{\mathbf{D}'}\mathbb{F}'$ .

3. If  $(G, \Psi, \psi)$  is a monoidal functor and  $F$  is left adjoint to  $G$  with unit  $\eta$ , then  $(F, \Phi, \phi)$  is opmonoidal, where  $\phi$  corresponds by adjunction to  $\psi$  and  $\Phi_{C,D}$  corresponds by adjunction to  $\Psi_{FC,FD} \circ (\eta_C \otimes \eta_D)$ . This defines a bijection between monoidal structures on  $G$  and opmonoidal structures on  $F$ .
4. If  $(G, \Psi, \psi)$  and  $(F, \Phi, \phi)$  are monoidal functors where  $F$  is left adjoint to  $G$ , then this adjunction is called a *monoidal adjunction*, provided that its unit and counit are monoidal transformations. In this situation, we have as follows:

- (a)  $(F, \Phi, \phi)$  is a strong monoidal and, hence, an opmonoidal functor;
- (b) The natural isomorphism  $\text{hom}(F-, -) \simeq \text{hom}(-, G-)$  is a monoidal isomorphism.

5. If  $(G, \Psi, \psi)$  is a monoidal and  $(F, \Phi, \phi)$  a strong opmonoidal (hence monoidal) functor, where  $F$  is left adjoint to  $G$ , then (b) above implies that the adjunction is monoidal.

If  $\mathbf{C}$  is symmetric monoidal then the internal hom functor  $[-, -] : \mathbf{C}^{\text{op}} \otimes \mathbf{C} \rightarrow \mathbf{C}$  is monoidal (see [4]) and, thus, also each functor  $[-, X] : \mathbf{C}^{\text{op}} \rightarrow \mathbf{C}$  is monoidal; this will not necessarily be the case for a nonsymmetric  $\mathbf{C}$ . For an important fragment of this, see however Section 1.2 below.

**Proposition 4** ([2, Chap. 3]). *Let  $\mathbb{F} := (F, \Phi, \phi) : \mathbb{C} \rightarrow \mathbb{C}'$  be a monoidal functor.*

1.  $\tilde{\mathbb{F}}(M, m, e) = (FM, FM \otimes FM \xrightarrow{\Phi_{M,M}} F(M \otimes M) \xrightarrow{Fm} FM, I' \xrightarrow{\phi} FI \xrightarrow{Fe} FM)$  and  $\tilde{\mathbb{F}}f = Ff$  defines an induced functor  $\tilde{\mathbb{F}} : \mathbf{Mon}\mathbb{C} \rightarrow \mathbf{Mon}\mathbb{C}'$ , such that the diagram

$$\begin{array}{ccc}
 \mathbf{Mon}\mathbb{C} & \xrightarrow{\tilde{\mathbb{F}}} & \mathbf{Mon}\mathbb{C}' \\
 U_m \downarrow & & \downarrow U'_m \\
 \mathbb{C} & \xrightarrow{F} & \mathbb{C}'
 \end{array}$$

*commutes (with forgetful functors  $U_m$  and  $U'_m$ ).*

2. *In the same way, one obtains a functor  $\hat{\mathbb{F}} : \mathbf{Sgr}\mathbb{C} \rightarrow \mathbf{Sgr}\mathbb{C}'$ , such that the diagram*

$$\begin{array}{ccc}
 \mathbf{Mon}\mathbb{C} & \xrightarrow{\tilde{\mathbb{F}}} & \mathbf{Mon}\mathbb{C}' \\
 \downarrow V & & \downarrow V' \\
 \mathbf{Sgr}\mathbb{C} & \xrightarrow{\hat{\mathbb{F}}} & \mathbf{Sgr}\mathbb{C}' \\
 \downarrow U_s & & \downarrow U'_s \\
 \mathbb{C} & \xrightarrow{F} & \mathbb{C}'
 \end{array}$$

*commutes (with forgetful functors  $V, V', U_s,$  and  $U'_s$ ).*

*Given a monoidal adjunction  $\mathbb{F} \dashv \mathbb{G}$  with unit  $\eta : Id \rightarrow GF$  and counit  $\epsilon : FG \rightarrow Id$ , then the induced functors  $\tilde{\mathbb{F}}$  and  $\tilde{\mathbb{G}}$  form an adjunction with unit  $\eta'$  and counit  $\epsilon'$ , such that  $U_m\eta' = \eta$  and  $U'_m\epsilon' = \epsilon$ . Similarly for  $\hat{\mathbb{F}}$  and  $\hat{\mathbb{G}}$ .*

### 1.2. Dualization in Monoidal Closed Categories

Let  $\mathbb{C}$  be a monoidal bi-closed category. We call the functor  $[-, I]_l$ , introduced in Section 1.1 above, the *left dualization functor* of  $\mathbb{C}$ . Analogously, there is the *right dualization functor*  $[-, I]_r$ .

Note that, from a categorical perspective, we should rather call these functors *semi-dualization functors* following [12, Def. 4.6], where the notion of a *semidualizable* object in a monoidal category is introduced. The linear dual of an  $R$ -module  $M$  is a dual in the categorical sense only, if  $M$  is reflexive. But since our focus is on modules and bimodules, we prefer to omit the prefix ‘semi’ in this note.

**Proposition 5.** *For every monoidal bi-closed category  $\mathbb{C}$ , the left and the right dualization functors form a dual adjunction, i.e.,  $\mathbb{C} \xrightarrow{[-, I]_r} \mathbb{C}^{op}$  is left adjoint to  $\mathbb{C}^{op} \xrightarrow{[-, I]_l} \mathbb{C}$ .*

*Proof.* Compose  $\text{hom}_{\mathbb{C}}(I, \Pi'_{C,D})$  and the inverse of  $\text{hom}_{\mathbb{C}}(I, \Pi_{C,D})$  (see Section 1.1) in order to obtain the requested natural isomorphisms

$$\text{hom}_{\mathbb{C}}(C, [D, I]_r) \simeq \text{hom}_{\mathbb{C}}(C \otimes D, I) \simeq \text{hom}_{\mathbb{C}}(D, [C, I]_l). \quad \square$$

**Theorem 6.** For every monoidal left closed category  $\mathbb{C}$ , the functor  $[-, I]_l : \mathbb{C}^{op} \rightarrow \mathbb{C}^l$  is monoidal.

*Proof.* Suppressing the constraints, we denote, for  $\mathbb{C}$ -objects  $C, D$ , by  $\bar{\Phi}_{C,D}$  the  $\mathbb{C}$ -morphism

$$(C \otimes D) \otimes [D, I]_l \otimes [C, I]_l \xrightarrow{\text{id} \otimes \text{ev}_{D,I}^l \otimes \text{id}} C \otimes I \otimes [C, I]_l \xrightarrow{\rho_C \otimes \text{id}} C \otimes [C, I]_l \xrightarrow{\text{ev}_{C,I}^l} I.$$

This family of morphisms is obviously natural in  $C$  and  $D$ .

Denoting by  $\Phi_{C,D} : [D, I]_l \otimes [C, I]_l \rightarrow [C \otimes D, I]_l$  the morphism corresponding to  $\bar{\Phi}_{C,D}$  by adjunction, i.e., the (unique) morphism making the following diagram commute

$$\begin{array}{ccc}
 (C \otimes D) \otimes [C \otimes D, I]_l & \xrightarrow{\text{ev}_{C \otimes D, I}^l} & I \\
 \uparrow \text{id} \otimes \Phi_{C,D} & & \uparrow \text{ev}_{C,I}^l \\
 (C \otimes D) \otimes [D, I]_l \otimes [C, I]_l & \xrightarrow{\text{id} \otimes \text{ev}_{D,I}^l \otimes \text{id}} & C \otimes I \otimes [C, I]_l \\
 & & \uparrow \rho_C \otimes \text{id}
 \end{array} \tag{1}$$

one thus obtains a natural transformation  $\Phi_{C,D} : [C, I]_l \otimes^l [D, I]_l \rightarrow [C \otimes D, I]_l$ .

Denoting by  $\phi := j_I : I \rightarrow [I, I]_l$  the isomorphism corresponding by adjunction to  $I \otimes I \xrightarrow{\rho_I = \lambda_I} I$  (see Section 1.1),  $([-, I]_l, \Phi, \phi)$  is a monoidal functor.

To prove coherence, commutativity of the following diagrams is to be shown:

$$\begin{array}{ccc}
 I \otimes [C, I]_l & \xrightarrow{\phi \otimes [C, I]_l} & [I, I]_l \otimes [C, I]_l & \xrightarrow{\Phi_{C,I}} & [C \otimes I, I]_l \\
 & \searrow \lambda_{[C, I]_l} & & \swarrow [\rho_C^{-1}, I] & \\
 & & [C, I]_l & & 
 \end{array} \tag{2}$$

$$\begin{array}{ccc}
 [C, I]_l \otimes I & \xrightarrow{[C, I]_l \otimes \phi} & [C, I]_l \otimes [I, I]_l & \xrightarrow{\Phi_{I,C}} & [I \otimes C, I]_l \\
 & \searrow \rho_{[C, I]_l} & & \swarrow [\lambda_C^{-1}, I] & \\
 & & [C, I]_l & & 
 \end{array} \tag{3}$$






Concerning Diagram (4), we first observe that the following diagrams commute, where we put  $\alpha = \alpha_{C,*B,*A}$  and  $\bar{\alpha} = \alpha_{A,B,C}$ . Again the cells referring to previously defined data are marked respectively; associativities are only labelled by  $\simeq$ . The top cell of the second diagram commutes by coherence. One thus gets

$$ev_{A(BC)} \circ (A(BC)) \otimes (\Phi_{A,BC} \circ \Phi_{BC} * A \circ \alpha) = ev_{A(BC)} \circ (A(BC)) \otimes ([\bar{\alpha}^{-1}, I] \circ \Phi_{AB,C} \circ *C\Phi_{A,B}),$$

which is equivalent to the commutativity of Diagram (4):



□

Since  $\mathbb{C}$  is left closed if and only if  $\mathbb{C}'$  is right closed (with  $[-, I]_l$  in  $\mathbb{C}$  being the same as  $[-, I]_r$  in  $\mathbb{C}'$  — see above), one immediately gets the following corollary.

**Corollary 7.** *For every monoidal right closed category  $\mathbb{C}$ , the functor  $[-, I]_r : \mathbb{C}^{op} \rightarrow \mathbb{C}'$  is monoidal.*

The monoidal structure on  $[-, I]_r$  is given by the isomorphism  $\psi := i_I : I \rightarrow [I, I]_r$  corresponding by adjunction to  $I \otimes I \xrightarrow{\lambda_I = \rho_I} I$  (see again Section 1.1) and the natural transformation  $\Psi$  whose components are those morphisms making the following diagram commute:

$$\begin{array}{ccc}
 [D \otimes C, I]_r \otimes (D \otimes C) & \xrightarrow{ev_{D \otimes C, I}^r} & I \\
 \uparrow \Psi_{D, C} \otimes \text{id} & & \uparrow ev_{C, I}^r \\
 [C, I]_r \otimes [D, I]_r \otimes (D \otimes C) & \xrightarrow{\text{id} \otimes ev_{D, I}^r \otimes \text{id}} & [C, I]_r \otimes I \otimes C \\
 & & \uparrow \text{id} \otimes \lambda_C \\
 & & [C, I]_r \otimes C
 \end{array}$$

**Remarks 8.**

1. The opmonoidal structure  $(\Psi, \psi)$  on  $[-, I]_r$  just defined corresponds to the monoidal structure  $(\Phi, \phi)$  on  $[-, I]_l$  in the sense of Remark 3 (3).
2. The natural transformation  $\Psi$  is related to the natural transformations  $\Theta_{C, D}^r$  and  $\Pi^r$  of Section 1.1 as follows:

$$\Psi_{C, D} = [D, I]_r \otimes [C, I]_r \xrightarrow{\Theta_{C, [D, I]_r}^r} [C, [D, I]_r]_r \xrightarrow{\Pi_{C, D}^r} [C \otimes D, I]_r$$

Similarly for  $\Phi$ .

**1.3. Bimodules**

Throughout this section,  $R$  denotes a not necessarily commutative unital ring and  ${}_R\mathbf{Mod}_R$  the category of  $R$ - $R$ -bimodules. We note that this category is abelian.

By  ${}_R\mathbf{Mod}_R \xrightarrow{|\cdot|} \mathbf{Ab}$ , we denote the underlying functor from  ${}_R\mathbf{Mod}_R$  into the category of abelian groups.

**1.3.1. The Monoidal Category of  $R$ - $R$ -Bimodules.** The category  ${}_R\mathbf{Mod}_R$  carries a monoidal structure given by the tensor product  $- \otimes_R -$  over  $R$  and the bimodule  $R$  as unit object. The resulting monoidal category  ${}_R\mathbf{Mod}_R$  is a nonsymmetric monoidal category if  $R$  fails to be commutative, and it is monoidal bi-closed (see, e.g., [5]) as follows:

- $[M, -]_r$ , the right adjoint of  $- \otimes_R M$ , is given by  $[M, N]_r = \text{hom}_R(M, N)$ , the  $R$ - $R$ -bimodule of right  $R$ -linear maps;

- $[M, -]_l$ , the right adjoint of  $M \otimes_R -$ , is given by  $[M, N]_l = {}_R \text{hom}(M, N)$ , the  $R$ - $R$ -bimodule of left  $R$ -linear maps.

In particular, one has the maps  $R \xrightarrow{j_R} [R, R]_l = {}_R \text{hom}(R, R)$  and  $R \xrightarrow{i_R} [R, R]_r = \text{hom}_R(R, R)$  (see Section 1.1). In the sequel, we will refer to these isomorphisms, which are the maps sending the unit  $1 \in R$  to the identity  $\text{id}_R$ , as  $\phi_R$  and  $\psi_R$ , respectively.

For bimodules  $M$  and  $N$ , the canonical surjection  $|M| \otimes_{\mathbb{Z}} |N| \rightarrow |M \otimes_R N|$  will be denoted by  ${}_R \text{can}_{M,N}$ . The family of these surjections forms a natural transformation. Denoting by  $\mathbb{Z} \xrightarrow{\chi_R} R$  the unique unital ring homomorphism, one gets a monoidal functor

$$(| - |, {}_R \text{can}, \chi_R): {}_R \mathbb{M} \text{od}_R \rightarrow \mathbb{A}b \tag{6}$$

from the monoidal category of bimodules into the monoidal category of abelian groups, the latter equipped with its standard tensor product.

With  $\tau$  the symmetry of the tensor product of abelian groups, one also gets a monoidal functor

$$(| - |, {}_R \text{can} \circ \tau, \chi_R): {}_R \mathbb{M} \text{od}_R^t \rightarrow \mathbb{A}b. \tag{7}$$

**1.3.2. A Monoidal Isomorphism.** With  $R^{\text{op}}$  the opposite ring of  $R$  one has a canonical isomorphism  ${}_R \mathbf{Mod}_R \rightarrow {}_{R^{\text{op}}} \mathbf{Mod}_{R^{\text{op}}}$ , switching left and right  $R$ -actions, which will be denoted by  $\overline{(-)}$ .

Note that there are natural isomorphisms  $\overline{N} \otimes_{R^{\text{op}}} \overline{M} \xrightarrow{\sigma_{M,N}} \overline{M \otimes_R N}$  acting as  $y \otimes_{R^{\text{op}}} x \mapsto x \otimes_R y$  (see, e.g., [8, p. 132]). Since<sup>2</sup>  $\overline{R} = R^{\text{op}}$ ,

$$(\overline{(-)}, \sigma, \text{id}_{R^{\text{op}}}) : {}_R \mathbb{M} \text{od}_R^t \rightarrow {}_{R^{\text{op}}} \mathbb{M} \text{od}_{R^{\text{op}}} \tag{8}$$

is a (strong) monoidal functor (in fact a monoidal isomorphism)  ${}_R \mathbb{M} \text{od}_R^t \simeq {}_{R^{\text{op}}} \mathbb{M} \text{od}_{R^{\text{op}}}$  and, thus, lifts by Proposition 4 to functorial isomorphisms

$$\begin{aligned} \mathbf{Mon}({}_R \mathbb{M} \text{od}_R) &= \mathbf{Mon}({}_R \mathbb{M} \text{od}_R^t) \simeq \mathbf{Mon}({}_{R^{\text{op}}} \mathbb{M} \text{od}_{R^{\text{op}}}), \\ \mathbf{Comon}({}_R \mathbb{M} \text{od}_R) &= \mathbf{Comon}({}_R \mathbb{M} \text{od}_R^t) \simeq \mathbf{Comon}({}_{R^{\text{op}}} \mathbb{M} \text{od}_{R^{\text{op}}}). \end{aligned}$$

**1.3.3. Dualization Functors for  ${}_R \mathbf{Mod}_R$ .** Recall from Section 1.2 the right and the left dualization functor on  ${}_R \mathbf{Mod}_R$ . These are the contravariant functors  $(-)^* := [-, R]_r$  and  $*(-) := [-, R]_l$ , which—on the level of sets—act as  $\text{hom}_R(-, R)$  and  ${}_R \text{hom}(-, R)$ , respectively (see Section 1.3.1).

Proposition 5 now specializes to the following proposition.

<sup>2</sup>Here  $R$  and  $R^{\text{op}}$  are considered as an  $R$ - $R$ -bimodule and an  $R^{\text{op}}$ - $R^{\text{op}}$ -bimodule, respectively.

**Proposition 9.** *Considering the left and right  $R$ -dualization functors as functors  $({}_R\mathbf{Mod}_R)^{op} \xrightarrow{*(-)} {}_R\mathbf{Mod}_R \xrightarrow{(-)^*} ({}_R\mathbf{Mod}_R)^{op}$ , the functor  $(-)^*$  is left adjoint to  $*(-)$ . The units and counits of this adjunction are the maps  $M \xrightarrow{\eta_M} *(M^*)$  and  $M \xrightarrow{\epsilon_M} (*M)^*$  with  $x \mapsto (y \mapsto y(x))$ .*

By Section 1.2, we obtain the following proposition.

**Proposition 10.**

1. *The dualization functors are monoidal functors as follows:*

- (a)  $\mathbb{D}_l := (*(-), \Phi, \phi_R) : {}_R\mathbb{M}od_R^{op} \rightarrow {}_R\mathbb{M}od_R^t ;$
- (b)  $\mathbb{D}_r := ((-)^*, \Psi, \psi_R) : {}_R\mathbb{M}od_R^{op} \rightarrow {}_R\mathbb{M}od_R^t .$

2. *Composing these with the strong monoidal functor of 1.3.2, one obtains the following monoidal functors:*

- (a)  $\overline{\mathbb{D}}_l := (\overline{*(-)}, \overline{\Phi} \circ \sigma, \overline{\phi_R}) : {}_R\mathbb{M}od_R^{op} \rightarrow {}_{R^{op}}\mathbb{M}od_{R^{op}} ;$
- (b)  $\overline{\mathbb{D}}_r := (\overline{(-)^*}, \overline{\Psi} \circ \sigma, \overline{\psi_R}) : {}_R\mathbb{M}od_R^{op} \rightarrow {}_{R^{op}}\mathbb{M}od_{R^{op}} .$

**Remark 11.** The action of the maps

$$*M \otimes_R *N \xrightarrow{\Phi_{N,M}} *(N \otimes_R M) \quad \text{and} \quad M^* \otimes_R N^* \xrightarrow{\Psi_{N,M}} (N \otimes_R M)^*$$

can, in view of the commutative diagrams of Section 1.2, be described by the following equations, for all  $\mu \in *M$  (resp.  $M^*$ ),  $v \in *N$  (resp.  $N^*$ ), and  $x \in M, y \in N$ :

$$(\Psi_{N,M}(\mu \otimes_R v))(y \otimes x) = \mu(v(y)x), \tag{9}$$

$$(\Phi_{N,M}(\mu \otimes_R v))(y \otimes x) = v(y\mu(x)). \tag{10}$$

Equivalently,

$$\Psi_{N,M}(\mu \otimes_R v) = \mu \circ \lambda_M \circ (v \otimes_R M), \tag{11}$$

$$\Phi_{N,M}(\mu \otimes_R v) = v \circ \rho_N \circ (N \otimes_R \mu). \tag{12}$$

Note that Sweedler’s natural transformation  $\mu$  of [14, Lemma 3.4 (b)], if specialized to  $X = Y = V = R$  and with the replacements  $R = M$  and  $S = N$  coincides with  $\Phi$ , while the natural transformation  $\eta$  of [14, Lemma 3.4 (a)] with the corresponding specializations and replacements coincides with  $\Psi$ .

We finally remark that the contravariant hom-functor  $M \mapsto {}_R\mathbf{Mod}_R(M, R)$ , from  ${}_R\mathbf{Mod}_R$  to  $\mathbf{Set}$ , can be lifted to a functor  $*(-)^* : {}_R\mathbf{Mod}_R^{op} \rightarrow \mathbf{Ab}$ . This functor can be supplied with a monoidal structure as follows: Denote by  $\Xi_{M,N} : M^* \otimes_{\mathbb{Z}} *N^* \rightarrow *(M \otimes_R N)^*$  the group homomorphism with  $\mu \otimes_{\mathbb{Z}} v \mapsto \rho_R \circ (\mu \otimes_R v)$ , by  $\mathbb{Z} \xrightarrow{\zeta} *R^*$  the group homomorphism with  $1 \mapsto \text{id}_R$ , and by  $\Xi'_{M,N}$  the composition  $*M^* \otimes_{\mathbb{Z}} *N^* \xrightarrow{\tau_{*M^*, *N^*}} *N^* \otimes_{\mathbb{Z}} *M^* \xrightarrow{\Xi_{N,M}} *(N \otimes_R M)^* = *(M \otimes'_R N)^*$ . Note

that this construction can be obtained by a purely categorical argument as well, using the fact that  ${}_R\mathbf{Mod}_R$  is enriched over  $\mathbf{Ab}$  (see [4]). Then the following proposition holds.

**Proposition 12.**

1.  $\mathbb{D} := (*(-)^*, \Xi, \xi) : {}_R\mathbb{M}od_R^{op} \rightarrow {}_R\mathbb{M}od_R^t \simeq {}_{R^{op}}\mathbb{M}od_{R^{op}}$  is a monoidal functor.
2.  $\mathbb{D}' := (*(-)^*, \Xi', \zeta) : ({}_R\mathbb{M}od_R^t)^{op} \rightarrow {}_R\mathbb{M}od_R^t \simeq {}_{R^{op}}\mathbb{M}od_{R^{op}}$  is a monoidal functor.

It is well known that every adjunction induces a largest equivalence. In more detail, let  $\mathbf{A} \xrightarrow{G} \mathbf{B} \xrightarrow{F} \mathbf{A}$  form an adjunction with unit  $\eta : \mathbf{Id}_{\mathbf{B}} \Rightarrow GF$  and counit  $\epsilon : FG \Rightarrow \mathbf{Id}_{\mathbf{A}}$ ; then the restrictions of  $G$  and  $F$  to the full subcategories  $Fix\eta$  and  $Fix\epsilon$  spanned by those objects of  $\mathbf{B}$  and  $\mathbf{A}$ , whose components of  $\eta$  and  $\epsilon$ , respectively, are isomorphisms, provide an equivalence. Also  $Fix\eta$  and  $Fix\epsilon$  obviously are the largest subcategories of  $\mathbf{B}$  and  $\mathbf{A}$ , respectively, which are equivalent under  $F$  and  $G$ . Applied to the adjunction of Proposition 9 above, we thus obtain a duality (= dual equivalence) between the full subcategories spanned by those bimodules  $M$  which are reflexive as left and right  $R$ -modules, respectively.

Characterizing these is not possible in general. However, one can describe a duality between somewhat smaller subcategories, induced by the dual adjunction  $(-)^* \dashv *(-)$ , which then even is a *monoidal* duality as follows, where we denote by  $\mathbf{FGP}_R$  and  ${}_R\mathbf{FGP}$  the full subcategories of  ${}_R\mathbf{Mod}_R$ , spanned by all bimodules which are finitely generated projective as right and as left  $R$ -modules, respectively (see also [14] and [15]).

**Proposition 13.**

1. The categories  $\mathbf{FGP}_R$  and  ${}_R\mathbf{FGP}$  are closed in  ${}_R\mathbf{Mod}_R$  under tensor products. In particular, these categories form monoidal subcategories  ${}_R\mathbf{FGP}$  and  $\mathbf{FGP}_R$  of  ${}_R\mathbb{M}od_R$ .
2. The dualization functors can be restricted as follows:

$$*(-) : {}_R\mathbf{FGP}^{op} \rightarrow \mathbf{FGP}_R \quad \text{and} \quad (-)^* : \mathbf{FGP}_R \rightarrow {}_R\mathbf{FGP}^{op},$$

and these restrictions provide a duality.

3. The restrictions of the monoidal functors  $\mathbb{D}_l = (*(-), \Phi, \phi_R)$  and  $\mathbb{D}_r = ((-)^*, \Psi, \psi_R)$ , considered as monoidal functors<sup>3</sup>,

$${}_R\mathbf{FGP}^{op} \xrightarrow{\mathbb{D}'_l} \mathbf{FGP}'_R \quad \text{and} \quad \mathbf{FGP}'_R \xrightarrow{\mathbb{D}'_r} {}_R\mathbf{FGP}^{op}$$

provide a monoidal duality.

4. Considering the dualization functors in the form of Proposition 10 (2) this amounts to a monoidal duality  ${}_R\mathbf{FGP}^{op} \simeq {}_{R^{op}}\mathbf{FGP}$ , given by  $\overline{\mathbb{D}'_l}$  and  $\overline{\mathbb{D}'_r}$ .

<sup>3</sup>Note that  $\Psi$  and  $\psi$  are invertible and, thus, provide the respective morphisms in  ${}_R\mathbf{FGP}^{op}$  by considering their inverses.

*Proof.*  $P$  and  $Q$  are  $R$ - $R$ -bimodules from  $\mathbf{FGP}_R$ , if and only if there exist retractions  $R^n \xrightarrow{p} P$  and  $R^m \xrightarrow{q} Q$  in  $\mathbf{Mod}_R$ . Then  $R^{nm} \simeq R^n \otimes_R R^m \xrightarrow{p \otimes_R q} P \otimes_R Q$  is a retraction in  $\mathbf{Mod}_R$ , such that  $P \otimes_R Q$  is finitely generated projective in  $\mathbf{Mod}_R$ . This proves 1, since the tensor product in  ${}_R\mathbf{Mod}_R$  has this module as its underlying right  $R$ -module.

To prove 2, it suffices to observe that (a)  $*R \simeq R$ , (b)  $*(-)$ , being a right adjoint by Proposition 9, preserves products and, thus, finite coproducts as well, since these are biproducts, and (c)  $*(-)$  preserves retracts by functoriality.

For 3, we first of all need to show that the restricted adjunction is a monoidal one, which (see Remark 3) is the case if and only if  $\Psi_{M,N}$  is an isomorphism, provided that  $M$  and  $N$  are finitely generated projective as right  $R$ -modules. By Remark 8, we have  $\Psi_{N,M} = \Pi'_{N,M} \circ \Theta_{N,M^*}$ , where  $\Pi'_{N,M}$  is an isomorphism; now  $\Theta_{N,M^*}$  is an isomorphism, provided that  $M$  or  $N$  is finitely generated projective (see, e.g., [1, 20. Ex. 12]). Consequently, the restriction of the dual adjunction to  ${}_R\mathbf{FGP}$  is monoidal and then, by 2, a monoidal duality as requested.  $\square$

## 2. $R$ -RINGS AND $R$ -CORINGS

### 2.1. The Categories of $R$ -Rings and $R$ -Corings

Recall that the category of monoids in  ${}_R\mathbf{Mod}_R^t \simeq {}_{R^{\text{op}}}\mathbf{Mod}_R^{\text{op}}$  is  ${}_1\mathbf{Ring}$ , the category of unital rings.

The *category of unital  $R$ -rings* is defined as  ${}_1\mathbf{Ring}_R := \mathbf{Mon}({}_R\mathbf{Mod}_R)$ , the category of monoids in  ${}_R\mathbf{Mod}_R$  (see [6, 3.24]) or as the comma category  $R \downarrow {}_1\mathbf{Ring}$  of unital rings under  $R$  (see [3]). In fact, these categories are easily seen to be isomorphic (see below). We will, in the sequel, use the first mentioned definition since it allows to make use of the theory of monoidal categories. (In particular, we can define the category  $\mathbf{Ring}_R$  of not necessarily unital  $R$ -rings as the category of semigroups in  ${}_R\mathbf{Mod}_R$ .)

However, when doing so one has to be careful, since one needs to distinguish  $R$ -rings and  $R^{\text{op}}$ -rings. As we will see below this distinction cannot be made in a simple categorical way: The categories  ${}_1\mathbf{Ring}_R$  and  ${}_1\mathbf{Ring}_{R^{\text{op}}}$  are isomorphic! Since  $\mathbf{Mon}({}_R\mathbf{Mod}_R) = \mathbf{Mon}({}_R\mathbf{Mod}_R')$ , they even may — or may not — be concretely isomorphic over  ${}_1\mathbf{Ring}$ , depending of the chosen forgetful functors (see below). Thus, if one wants to distinguish them categorically, one only can do that on the level of concrete categories.

The category of comonoids in  ${}_R\mathbf{Mod}_R$  is called the *category of counital  $R$ -corings* (see [6]) and will be denoted by  ${}_\epsilon\mathbf{Coring}_R$ ;  $\mathbf{Coring}_R$  then is the category of co-semigroups in  ${}_R\mathbf{Mod}_R$ .

**The Underlying Ring of an  $R$ -Ring** The monoidal functor of Eq. (6) induces a functor  $I_{\otimes_R} : \mathbf{Mon}({}_R\mathbf{Mod}_R) \rightarrow {}_1\mathbf{Ring}$  (see Proposition 4), which obviously is faithful<sup>4</sup>:

$I_{\otimes_R}$  maps a monoid  $(M, m, e)$  to  $(M, M \otimes_{\mathbb{Z}} M \xrightarrow{R\text{can}} M \otimes_R M \xrightarrow{m} M, \mathbb{Z} \xrightarrow{\gamma_R} R \xrightarrow{e} M)$ , called the *underlying ring of  $(M, m, e)$* ; in particular,  $I_{\otimes_R}(R) = R$ .

<sup>4</sup>We use the somewhat clumsy notation  $I_{\otimes_R}$  in order to stress the fact that this functor depends on the monoidal structure.

Analogously there is a functor  $I_{\otimes_R}^t : \mathbf{Mon}(R\mathbf{Mod}_R^t) = \mathbf{Mon}(R\mathbf{Mod}_R) \rightarrow {}_1\mathbf{Ring}$ , which is faithful as well:

$I_{\otimes_R}^t$  maps a monoid  $(M, m, e)$  to  $(M, M \otimes_{\mathbb{Z}} M \xrightarrow{R\text{can}^{\text{op}}} M \otimes_R M \xrightarrow{m} M, \mathbb{Z} \xrightarrow{\chi_R} R \xrightarrow{e} M)$ . Hence,  $I_{\otimes_R}^t(M, m, e) = (I_{\otimes_R}(M, m, e))^{\text{op}}$ . In particular,  $I_{\otimes_R}^t(R) = R^{\text{op}}$ .

**2.1.1. Some Categorical Isomorphisms.** Let  $(M, m, e)$  be an  $R$ -ring. Since, as for every monoid in a monoidal category, the unit  $e$  is a monoid homomorphism, the morphism  $I_{\otimes_R} e = e$  is a ring homomorphism from  $R$  into the underlying ring of  $(M, m, e)$ . In other words,  $(I_{\otimes_R}(M, m, e), e)$  is an object of  $R \downarrow {}_1\mathbf{Ring}$ . This defines a functor  $\Phi_R : \mathbf{Mon}(R\mathbf{Mod}_R) \rightarrow R \downarrow {}_1\mathbf{Ring}$  if one puts  $\Phi_R f = f$  for each morphism  $f$  of monoids in  $R\mathbf{Mod}_R$ .

Conversely, if  $(M, p, u)$  is a monoid in  $R\mathbf{Mod}_R^t \simeq R^{\text{op}}\mathbf{Mod}_{R^{\text{op}}}$ , i.e., a unital ring and  $e : R \rightarrow (M, p, u)$  a morphism in  ${}_1\mathbf{Ring}$ ,  $M$  becomes an  $R$ - $R$ -bimodule in the obvious way. Then  $e$  is a morphism in  ${}_R\mathbf{Mod}_R$  and  $p$  induces a ( $R$ -left and -right linear) map  $m : M \otimes_R M \rightarrow M$  with  $m \circ R\text{can} = p$ . It is well known (and easy to see) that the assignment  $((M, p, u), e) \mapsto (M, m, e)$  defines a functor  $\Psi_R : R \downarrow {}_1\mathbf{Ring} \rightarrow \mathbf{Mon}(R\mathbf{Mod}_R)$ , if one puts  $\Psi_R f = f$ , and this is the inverse of  $\Phi_R$ .

Denoting by  $V_R$  the familiar forgetful functor from  $R \downarrow {}_1\mathbf{Ring}$  to  ${}_1\mathbf{Ring}$  the following diagram commutes:

$$\begin{array}{ccc}
 R \downarrow {}_1\mathbf{Ring} & \begin{array}{c} \xleftarrow{\Psi_R} \\ \xrightarrow{\Phi_R} \end{array} & \mathbf{Mon}(R\mathbf{Mod}_R) \\
 \searrow V_R & & \swarrow I_{\otimes_R} \\
 & {}_1\mathbf{Ring} &
 \end{array}$$

Thus, the categories  $R \downarrow {}_1\mathbf{Ring}$  and  $\mathbf{Mon}(R\mathbf{Mod}_R)$  are isomorphic as concrete categories over  ${}_1\mathbf{Ring}$ .

Since the categories  $\mathbf{Mon}(R\mathbf{Mod}_R)$  and  $\mathbf{Mon}(R\mathbf{Mod}_R^t)$  coincide, the monoidal isomorphism of Eq. (8) induces a functor  $\Omega : {}_1\mathbf{Ring}_R \rightarrow {}_1\mathbf{Ring}_{R^{\text{op}}}$ , and this is an isomorphism; it makes the following diagram commute:

$$\begin{array}{ccc}
 \mathbf{Mon}(R\mathbf{Mod}_R) & \xrightarrow{\Omega} & \mathbf{Mon}(R^{\text{op}}\mathbf{Mod}_{R^{\text{op}}}) \\
 \searrow I_{\otimes_R}^t & & \swarrow I_{\otimes_{R^{\text{op}}}} \\
 & {}_1\mathbf{Ring} &
 \end{array} \tag{13}$$

The isomorphism  $\Omega$  corresponds to the isomorphism  $\Sigma : R \downarrow {}_1\mathbf{Ring} \rightarrow R^{\text{op}} \downarrow {}_1\mathbf{Ring}$ , which maps a ring homomorphism  $R \xrightarrow{f} S$  to  $R^{\text{op}} \xrightarrow{f} S^{\text{op}}$ . In particular, the following diagram commutes:

$$\begin{array}{ccc}
 \mathbf{Mon}(R\mathbf{Mod}_R) & \xrightarrow{\Phi_R} & R \downarrow {}_1\mathbf{Ring} \\
 \Omega \downarrow & & \downarrow \Sigma \\
 \mathbf{Mon}(R^{\text{op}}\mathbf{Mod}_{R^{\text{op}}}) & \xrightarrow{\Phi_{R^{\text{op}}}} & R^{\text{op}} \downarrow {}_1\mathbf{Ring}
 \end{array}$$



Similarly, there is a functor  $\Omega' : \mathbf{Sgr}({}_R\mathbb{M}\text{od}_R) \rightarrow \mathbf{Sgr}({}_{R^{\text{op}}}\mathbb{M}\text{od}_{R^{\text{op}}})$ .

Summarizing this, we state as follows: There are the following isomorphisms of concrete categories over  ${}_1\mathbf{Ring}$ , characterizing the concrete categories of  $R$ -rings and  $R^{\text{op}}$ -rings, respectively:

1.  $(R \downarrow {}_1\mathbf{Ring}, V_R) \xrightarrow{\Psi_R} (\mathbf{Mon}({}_R\mathbb{M}\text{od}_R), I_{\otimes_R})$ ;
2.  $(R^{\text{op}} \downarrow {}_1\mathbf{Ring}, V_{R^{\text{op}}}) \xrightarrow{\Psi_{R^{\text{op}}}} (\mathbf{Mon}({}_{R^{\text{op}}}\mathbb{M}\text{od}_{R^{\text{op}}}), I_{\otimes_{R^{\text{op}}}}) \xrightarrow{\Omega^{-1}} (\mathbf{Mon}({}_R\mathbb{M}\text{od}_R^t), I_{\otimes_R^t}) = (\mathbf{Mon}({}_R\mathbb{M}\text{od}_R), I_{\otimes_R^t})$ .

### 2.2. Dual Rings of a Coring

**2.2.1. Dual Ring Functors.** Using the equation  ${}_{\epsilon}\mathbf{Coring}_R^{\text{op}} = (\mathbf{Comon}({}_R\mathbb{M}\text{od}_R))^{\text{op}} = \mathbf{Mon}({}_{(R\mathbb{M}\text{od}_R)^{\text{op}}})$ , the left and right monoidal dualization functors  $\mathbb{D}_l$  and  $\mathbb{D}_r$  induce, by Proposition 4, functors

$$\widetilde{\mathbb{D}}_l, \widetilde{\mathbb{D}}_r : {}_{\epsilon}\mathbf{Coring}_R^{\text{op}} \rightarrow {}_1\mathbf{Ring}_R \quad \text{and} \quad \widehat{\mathbb{D}}_l, \widehat{\mathbb{D}}_r : \mathbf{Coring}_R^{\text{op}} \rightarrow \mathbf{Ring}_R,$$

while the left and right monoidal dualization functors  $\overline{\mathbb{D}}_l$  and  $\overline{\mathbb{D}}_r$  induce functors

$$\widetilde{\overline{\mathbb{D}}}_l, \widetilde{\overline{\mathbb{D}}}_r : {}_{\epsilon}\mathbf{Coring}_R^{\text{op}} \rightarrow {}_1\mathbf{Ring}_{R^{\text{op}}} \quad \text{and} \quad \widehat{\overline{\mathbb{D}}}_l, \widehat{\overline{\mathbb{D}}}_r : \mathbf{Coring}_R^{\text{op}} \rightarrow \mathbf{Ring}_{R^{\text{op}}}$$

such that (for the left dualization functors) the following diagram commutes, where  $|-|$  denotes the various forgetful functors:

$$\begin{array}{ccccc}
 & & \widetilde{\overline{\mathbb{D}}}_l & & \\
 & & \curvearrowright & & \\
 & & \widetilde{\mathbb{D}}_l & & \\
 & & \curvearrowleft & & \\
 & & \Omega & & \\
 {}_{\epsilon}\mathbf{Coring}_R^{\text{op}} & \xrightarrow{\widetilde{\mathbb{D}}_l} & {}_1\mathbf{Ring}_R & \xrightarrow{\Omega} & {}_1\mathbf{Ring}_{R^{\text{op}}} & (14) \\
 \downarrow V' & & \downarrow V & & \downarrow \bar{V} \\
 \text{Coring}_R^{\text{op}} & \xrightarrow{\widehat{\mathbb{D}}_l} & \mathbf{Ring}_R & \xrightarrow{\Omega'} & \mathbf{Ring}_{R^{\text{op}}} \\
 \downarrow |-| & & \downarrow & & \downarrow |-| \\
 {}_R\mathbf{Mod}_{R^{\text{op}}} & \xrightarrow{*(-)} & {}_R\mathbf{Mod}_R & \xrightarrow{\overline{(-)}} & {}_{R^{\text{op}}}\mathbf{Mod}_{R^{\text{op}}} \\
 \downarrow U_l^{\text{op}} & & \downarrow & & \downarrow |-| \\
 {}_R\mathbf{Mod}^{\text{op}} & \xrightarrow{{}_R\text{hom}(-, R)} & & & \mathbf{Set} .
 \end{array}$$

In the corresponding diagram for the right dualization functors, the lower cell is


$$\begin{array}{ccccc}
 {}_R\mathbf{Mod}_R^{\text{op}} & \xrightarrow{(-)^*} & {}_R\mathbf{Mod}_R & \xrightarrow{\overline{(-)}} & {}_{R^{\text{op}}}\mathbf{Mod}_{R^{\text{op}}} & (15) \\
 \downarrow U_r^{\text{op}} & & & & \downarrow |-| \\
 \mathbf{Mod}_R^{\text{op}} & \xrightarrow{\text{hom}_R(-, R)} & & & \mathbf{Set} .
 \end{array}$$

**Remarks 14.**

1. For the sake of clarity, we stress the fact already mentioned above that it can be seen as a matter of taste whether the dual ring of an  $R$ -coring is an  $R$ -ring or an  $R^{\text{op}}$ -ring! Both options are available and they correspond to each other by the functor  $\Omega$ . Both, the more natural (using “natural” tensor products only) and the more conceptual (consider the dual ring functor as a functor induced by a monoidal functor) point of view, suggest however as follows: *The dual ring of an  $R$ -coring is an  $R^{\text{op}}$ -ring.*
2. Clearly the monoidal functors  $\mathbb{D}$  and  $\mathbb{D}'$  from Proposition 12 induce, again by Proposition 4, functors  $\widetilde{\mathbb{D}}, \widetilde{\mathbb{D}}' : {}_\epsilon\mathbf{Coring}_R^{\text{op}} \rightarrow {}_1\mathbf{Ring}$  (recall that  $\mathbf{Comon}({}_R\mathbb{M}\circledast R) = \mathbf{Comon}({}_R\mathbb{M}\circledast R) = {}_\epsilon\mathbf{Coring}_R$  by Fact 1).

**2.2.2. The Ring Structure of Dual Rings.** We next describe the various dual rings constructed above as objects of  ${}_1\mathbf{Ring} \downarrow R, {}_1\mathbf{Ring} \downarrow R^{\text{op}}$ , and  ${}_1\mathbf{Ring}$ , respectively, i.e., in the form  $(I(X, m, e), e)$ , where  $I$  is the respective forgetful functor and  $I(X, m, e)$  is a unital ring (considered as a monoid in  $\mathbb{A}\mathbb{b}$ ) whose unit  $e$  (considered as a group homomorphism  $R \rightarrow X$ ). Then in fact is a unital ring homomorphism  $I(R) \rightarrow I(X, m, e)$ . This enables us in particular to compare our constructions with those of [14] as follows.

Given any  $R$ -coring  $(C, \Delta, \epsilon)$ , we note first that, by Section 2.1, the underlying rings of the  $R^{\text{op}}$ -ring  $\widetilde{\mathbb{D}}_l(C, \Delta, \epsilon)$  and  $\widetilde{\mathbb{D}}_r(C, \Delta, \epsilon)$  are the opposite rings of the underlying rings of the  $R$ -rings  $\mathbb{D}_l(C, \Delta, \epsilon)$  and  $\mathbb{D}_r(C, \Delta, \epsilon)$ , respectively. The situation is illustrated by the following commutative diagram:



It thus suffices to consider the latter ones.

Since  ${}^*\Delta$  (and  $\Delta^*$  and  ${}^*\Delta^*$  as well) acts by pre-composition with  $\Delta$  we get, by definition of  $\Phi$ ,  $\Psi$ , and  $\Xi$ , respectively, the following descriptions (where  $(\mu, \nu) \in {}^*C \times {}^*C$  (resp.  $C^* \times C^*$ ):

1.  $\widetilde{\mathbb{D}}_l(C, \Delta, \epsilon)$  is the unital  $R$ -ring  $({}^*C, m_l, e_l)$  with

$$m_l = {}^*C \otimes_R^l {}^*C \xrightarrow{\Phi_{C,C}} {}^*(C \otimes_R C) \xrightarrow{{}^*\Delta} {}^*C \quad \text{and} \quad e_l = R \xrightarrow{\phi_R} {}^*R \xrightarrow{{}^*\epsilon} {}^*C.$$

(a) Its underlying unital ring thus is

$$I_{\otimes_R}(\widetilde{\mathbb{D}}_l(C, \Delta, \epsilon)) = ({}_R\text{hom}(C, R), m_l \circ {}_R\text{can}, e_l \circ \chi_R)$$

with multiplication

$$\begin{aligned} \mu \cdot \nu &= m_l(\mu \otimes_R \nu) = \Phi_{C,C}(\mu \otimes_R \nu) \circ \Delta \\ &= C \xrightarrow{\Delta} C \otimes_R C \xrightarrow{C \otimes \mu} C \otimes_R R \xrightarrow{\rho_C} C \xrightarrow{\nu} R \end{aligned}$$

and unital element  $u_l = e_l \circ \chi_R(1) = {}^*\epsilon \circ \phi_R \circ \chi_R(1) = {}^*\epsilon(\text{id}_R) = \epsilon$ .

(b) The map  $e_l$  acts as  $e_l(r) = \epsilon(cr)$ , for  $c \in C$  and  $r \in R$ .

2.  $\widetilde{\mathbb{D}}_r(C, \Delta, \epsilon)$  is the unital  $R$ -ring  $(C^*, m_r, e_r)$  with

$$m_r = C^* \otimes_R^l C^* \xrightarrow{\Psi_{C,C}} (C \otimes_R C)^* \xrightarrow{\Delta^*} C^* \quad \text{and} \quad e_r = R \xrightarrow{\psi_R} R^* \xrightarrow{\epsilon^*} C^*.$$

(a) Its underlying unital ring thus is

$$I_{\otimes_R}(\widetilde{\mathbb{D}}_r(C, \Delta, \epsilon)) = (\text{hom}_R(C, R), m_r \circ {}_R\text{can}, e_r \circ \chi_R)$$

with multiplication

$$\begin{aligned} \mu \cdot \nu &= m_r(\mu \otimes_R \nu) = \Psi_{C,C}(\mu \otimes_R \nu) \circ \Delta \\ &= C \xrightarrow{\Delta} C \otimes_R C \xrightarrow{\nu \otimes C} C \otimes_R R \xrightarrow{\lambda_C} C \xrightarrow{\mu} R \end{aligned}$$

and unital element  $u_r = e_r \circ \chi_R(1) = \epsilon^* \circ \psi_R \circ \chi_R(1) = \epsilon^*(\text{id}_R) = \epsilon$ .

(b) The map  $e_r$  acts as  $e_r(r)(c) = \epsilon(rc)$ , for  $c \in C$  and  $r \in R$ .

3.  $\widetilde{\mathbb{D}}(C, \Delta, \epsilon)$  is the unital ring  $({}^*C^*, m, e)$  with

$$m = {}^*C^* \otimes_{\mathbb{Z}} {}^*C^* \xrightarrow{\Xi_{C,C}} {}^*(C \otimes_R C)^* \xrightarrow{{}^*\Delta^*} {}^*C^* \quad \text{and} \quad e = \mathbb{Z} \xrightarrow{\zeta} {}^*R^* \xrightarrow{{}^*\epsilon^*} {}^*C^*.$$

Its multiplication, thus, is given by

$$\begin{aligned} \mu \cdot \nu &= m(\mu \otimes_{\mathbb{Z}} \nu) = \Xi_{C,C}(\mu \otimes_{\mathbb{Z}} \nu) \circ \Delta \\ &= C \xrightarrow{\Delta} C \otimes_R C \xrightarrow{\mu \otimes \nu} R \otimes_R R \xrightarrow{\rho_R} R, \end{aligned}$$

while its unital element is  $\epsilon$ .

4.  $\widetilde{\mathbb{D}}'(C, \Delta, \epsilon) = (\widetilde{\mathbb{D}}(C, \Delta, \epsilon))^{\text{op}}$ .

**Remark 15.**

1. When comparing our constructions with Sweedler’s (see Introduction), one gets the following equations by inspection:

$$(a) \text{Sw}_l(C, \Delta, \epsilon) = (I_{\otimes_R}(\widetilde{\mathbb{D}}_l(C, \Delta, \epsilon)))^{\text{op}} = I_{\otimes_R^l}(\widetilde{\mathbb{D}}_l(C, \Delta, \epsilon)) = I_{\otimes_{R^{\text{op}}}}(\widetilde{\mathbb{D}}_l(C, \Delta, \epsilon));$$

$$(b) \lambda_C = e_l.$$

Consequently, up to the isomorphism  $\Phi_{R^{\text{op}}}$  of Section 2, Sweedler’s construction is nothing but the functor  $\widetilde{\mathbb{D}}_l$ , which clearly coincides with Takeuchi’s functor  $\mathbb{D}_l$ .

2. Similarly, Sweedler’s construction  $(\text{Sw}_r(C, \Delta, \epsilon), \rho_C)$  coincides with  $\widetilde{\mathbb{D}}_r$ , up to the isomorphism  $\Phi_{R^{\text{op}}}$ .
3. In Example 3.6 [14] Sweedler provides a multiplication on  $C^* = \text{hom}_R(C, R)$ , which—since, what there is called  $\eta$  is our  $\Psi_{C,C}$  (see Remark 11)—is the map  $m_r$  of 2. above. The ring he describes here is  $I_{\otimes_R}(\widetilde{\mathbb{D}}_l(C, \Delta, \epsilon))$ . The difference, thus, can be described as follows: while in his original construction he applies the underlying functor  $I_{\otimes_R}$  to the monoidal construction  $\widetilde{\mathbb{D}}_l(C, \Delta, \epsilon)$ , here he applies the underlying functor  $I_{\otimes_R^l}$ ; and  $I_{\otimes_R^l}(M, m, e)$  equals  $(I_{\otimes_R}(M, m, e))^{\text{op}}$  (see above).
4. The following result should be expected and is easy to check: Take an  $R$ -coring  $(C, \Delta, \epsilon)$  and form the dual  $R^{\text{op}}$ -rings  $\widetilde{\mathbb{D}}_l(C, \Delta, \epsilon)$  and  $\widetilde{\mathbb{D}}_r(C, \Delta, \epsilon)$ . Now consider  $(C, \Delta, \epsilon)$  as an  $R^{\text{op}}$ -coring and form the respective dual  $R$ -rings, using the monoidal isomorphism from Section 1.3.2, which allows for seeing an  $R$ -(co)ring as an  $R^{\text{op}}$ -(co)ring. Then the left dual  $R$ -ring of  $(C, \Delta, \epsilon)$  coincides with the right dual  $R^{\text{op}}$ -ring  $\widetilde{\mathbb{D}}_r(C, \Delta, \epsilon)$ , considered as an  $R$ -ring.
5. We leave it to the reader to state the respective statements concerning not necessarily unital dual rings of not necessarily counital  $R$ -corings.

**Remark 16.** If  $R$  is a commutative ring and  $\mathbf{Mod}_R$  is considered instead of  ${}_R\mathbf{Mod}_R$ , all of these construction coincide, as is immediate from the commutativity of

$$\begin{array}{ccccc}
 & & \mu \otimes_R \nu & & \mu \otimes_R \nu \\
 & \swarrow & & \searrow & \\
 C \otimes_R C & \xrightarrow{C \otimes_R \nu} & C \otimes_R R & \xrightarrow{\mu \otimes_R R} & R \otimes_R R & \xleftarrow{R \otimes_R \nu} & R \otimes_R C & \xleftarrow{\mu \otimes_R C} & C \otimes_R C \\
 & \downarrow \rho_C & & \downarrow \rho_R = \lambda_R & & \downarrow \lambda_C & & & \\
 & C & \xrightarrow{\mu} & R & \xleftarrow{\nu} & C & & & 
 \end{array}$$

which is a consequence of the fact that, for  $R$  commutative, the monoidal structure on  $\mathbf{Mod}_R$  is symmetric.

Moreover, for each  $R$ -coalgebra  $\mathbf{C}$  this construction not only gives a monoid in  ${}_R\mathbf{Mod}_R^l \simeq {}_{R^{\text{op}}}\mathbf{Mod}_{R^{\text{op}}}$  but even a monoid in  $\mathbb{A}b$ , i.e., an  $R$ -algebra, the so-called *dual algebra of the coalgebra  $\mathbf{C}$* . This in fact is a special instance of the familiar *convolution algebra* construction. The latter is best described as follows (see e.g. [4]):

1. For each symmetric monoidal closed category  $\mathbf{C}$  its internal hom-functor is a monoidal functor  $[-, -] : \mathbf{C}^{\text{op}} \otimes \mathbf{C} \rightarrow \mathbf{C}$ .

2. Noting that  $\mathbf{Mon}(\mathbb{C}^{\text{op}} \otimes \mathbb{C}) = \mathbf{Mon}\mathbb{C}^{\text{op}} \times \mathbf{Mon}\mathbb{C} = (\mathbf{Comon}\mathbb{C})^{\text{op}} \times \mathbf{Mon}\mathbb{C}$  one thus obtains a functor  $[-, -] : (\mathbf{Comon}\mathbb{C})^{\text{op}} \times \mathbf{Mon}\mathbb{C} \rightarrow \mathbf{Mon}\mathbb{C}$  which, in case  $\mathbb{C} = \mathbb{A}b$ , is the convolution algebra functor.

Proposition 13 then takes the form of the generalization of the familiar duality between the categories  $\mathbf{Coalg}_k$  of  $k$ -algebras and  $\mathbf{Alg}_k$  of  $k$ -algebras for a field  $k$  (see e.g. [7]) to arbitrary commutative rings: the categories  ${}_{fgp}\mathbf{Coalg}_R$  and  ${}_{fgp}\mathbf{Alg}_R$  of  $R$ -coalgebras and  $R$ -algebras, respectively, which are finitely generated projective as  $R$ -modules, are dually equivalent.

### 2.3. Dual Corings of $R$ -Rings

By Proposition 13, one can restrict the functors  $\mathbb{D}_l$  and  $\mathbb{D}_r$  to obtain a monoidal equivalence, given by the restrictions  $\widetilde{\mathbb{D}}'_l : {}_R\mathbf{FGIP}^{\text{op}} \rightarrow {}_{R^{\text{op}}}\mathbf{FGIP}$  and  $\widetilde{\mathbb{D}}'_r : {}_{R^{\text{op}}}\mathbf{FGIP} \rightarrow {}_R\mathbf{FGIP}^{\text{op}}$ . By Proposition 4, the induced functors

$$\begin{aligned} \widetilde{\mathbb{D}}'_l &: \mathbf{Mon}({}_R\mathbf{FGIP}^{\text{op}}) \rightarrow \mathbf{Mon}({}_{R^{\text{op}}}\mathbf{FGIP}) \quad \text{and} \\ \widetilde{\mathbb{D}}'_r &: \mathbf{Mon}({}_{R^{\text{op}}}\mathbf{FGIP}) \rightarrow \mathbf{Mon}({}_R\mathbf{FGIP}^{\text{op}}). \end{aligned}$$

form an adjunction, where the natural isomorphisms  $\text{Id} \Rightarrow \widetilde{\mathbb{D}}'_l \widetilde{\mathbb{D}}'_r$  and  $\widetilde{\mathbb{D}}'_r \widetilde{\mathbb{D}}'_l \Rightarrow \text{Id}$  coincide, in  ${}_R\mathbf{Mod}_R$ , with the units and counits of the adjunction  $(-)^* \dashv^* (-)$ . Since the latter are isomorphisms, this adjunction in fact is a duality.

In other words, one not only can assign to each (counital)  $R$ -coring  $\mathbb{C}$  with underlying module  $C$  in  ${}_R\mathbf{FGP}$  a dual (unital)  $R^{\text{op}}$ -ring (whose underlying module lies in  ${}_{R^{\text{op}}}\mathbf{FGP}$ ), but conversely, one can assign to each such coring a dual  $R$ -ring, and this defines a dual equivalence between the subcategory  $\mathbf{Comon}({}_R\mathbf{FGP})$  of  ${}_{\epsilon}\mathbf{Coring}_R$  and the subcategory  $\mathbf{Mon}({}_{R^{\text{op}}}\mathbf{FGP})$  of  ${}_1\mathbf{Ring}_{R^{\text{op}}}$ . Similarly, there is also a duality  $(\mathbf{Comon}(\mathbf{IFGP}_R))^{\text{op}} \simeq \mathbf{Mon}(\mathbf{IFGP}_{R^{\text{op}}})$ . This is the content of the *Dual Coring Theorem* of [14] (see also [15]).

Note that these dualities also exist in the non(co)unital case.

### 3. UNITARIZATION AND COUNITARIZATION

Returning to Diagram (14) we recall that, by Theorem 5 and Remark 6 of [11], the functors  $V$  and  $\bar{V}$  have left adjoints  $A$  and  $\bar{A}$  respectively, the *unitarization functors*;  $V'$ , the dual of the *counitarization functor*, has a left adjoint as well. Since  $*(-)$  is right adjoint to  $(-)^*$ ,  $*(-)$  preserves (binary) products and  $(-)^*$  preserves (binary) coproducts. Since  ${}_R\mathbf{Mod}_R$  has biproducts and both dualization functors are additive, they preserve binary products and binary coproducts. Thus, the hypothesis of Theorem 10 of [11] are satisfied by both dualization functors and we get the following compatibility results for the operations of (co)unitarization and (left and right) dualization.

**Proposition 17.** *Let  $R$  be a unital ring. Then the following hold for every (not necessarily counital)  $R$ -coring  $(C, \Delta)$ :*

1. *The unitarization of the left dual  $\widehat{\mathbb{D}}_l(C, \Delta)$  coincides with the left dual  $\widetilde{\mathbb{D}}_l A'(C, \Delta)$  of the counitarization of  $(C, \Delta)$ ;*

2. The unitarization of the right dual  $\widehat{\mathbb{D}}_r(C, \Delta)$  coincides with the right dual  $\widetilde{\mathbb{D}}_r A'(C, \Delta)$  of the counitarization of  $(C, \Delta)$ .

Analogous statements clearly hold with respect to the dual ring functors  $\widetilde{\mathbb{D}}_l$  and  $\widetilde{\mathbb{D}}_r$  and with respect to the functors  $\widetilde{\mathbb{D}}$  and  $\widetilde{\mathbb{D}}'$ . We leave the precise formulation to the reader.

## REFERENCES

- [1] Anderson, F. W., Fuller, K. R. (1974). *Rings and Categories of Modules*. New York–Heidelberg–Berlin: Springer.
- [2] Aguiar, M., Mahajan, S. (2010). *Monoidal Functors, Species and Hopf Algebras*. CRM Monograph Series, Vol. 29. Providence, RI: American Mathematical Society.
- [3] Bergman, G. M., Hausknecht, A. O. (1996). Cogroups and co-rings in categories of associative rings. *Memoirs of the American Mathematical Society* 45.
- [4] Booker, Th., Street, R. (2013). Tannaka duality and convolution for duoidal categories. *Theory and Applications of Categories* 28(6):166–205.
- [5] Borceux, F. (1994). Handbook of categorical algebra 2. *Encyclopedia of Mathematics and Its Applications*, vol. 51. Cambridge University Press.
- [6] Brzezinski, T. (2009). Comodules and corings. In: Hazewinkel, M., ed. *Handbook of Algebra*. Vol. 6. Amsterdam: Elsevier, pp. 237–318.
- [7] Dăscălescu, S. Năstăsescu, C., Raianu, Ș. (2001). *Hopf Algebras: An Introduction*. Pure and Applied Mathematics, Vol. 235. Marcel Dekker, Inc.
- [8] Jacobson, N. (1980). *Basic Algebra II*. San Francisco: W.H. Freeman and Company.
- [9] Kelly, G. M. (1982). *Basic Concepts of Enriched Category Theory*. Lecture Notes in Mathematics, Vol. 64. Cambridge University Press.
- [10] Mac Lane, S. (1998). *Categories for the Working Mathematician*. 2nd ed. New York: Springer.
- [11] Poinsot, L., Porst, H.-E. (2015). Free monoids over semigroups in a monoidal category: Construction and applications. *Comm. Algebra* 43:4873–4899.
- [12] Stolz, S., Teichner, P. (2012). Traces in monoidal categories. *Trans. Amer. Math. Soc.* 364:4425–4464.
- [13] Street, R. (2007). *Quantum Groups*. Cambridge: Cambridge University Press.
- [14] Sweedler, M. (1975). The predual theorem to the Jacobson-Bourbaki theorem. *Trans. Amer. Math. Soc.* 213:391–406.
- [15] Takeuchi, M. (1987).  $\sqrt{\text{Morita}}$  theory: Formal ring laws and monoidal equivalences of categories of bimodules. *J. Math. Soc. Japan* 39:301–336.