

The Redei–Berge symmetric function of a directed graph


Darij Grinberg (Drexel University)
joint work with Richard P. Stanley

Haverford College, 2023-11-06
CAP, 2023-11-15


slides: <http://www.cip.ifi.lmu.de/~grinberg/algebra/haverford2023.pdf>

paper (draft): <https://arxiv.org/abs/2307.05569>

- **Definition.** A **digraph** (= directed graph) means a pair (V, A) of a finite set V and a subset $A \subseteq V \times V$. The elements $(u, v) \in A$ are called **arcs** of this digraph, and are drawn accordingly.
- **Example.**



- **Definition.** A **digraph** (= directed graph) means a pair (V, A) of a finite set V and a subset $A \subseteq V \times V$. The elements $(u, v) \in A$ are called **arcs** of this digraph, and are drawn accordingly.
- **Example.**




- Thus, we allow loops $((u, u) \in A)$ and antiparallel arcs $((u, v) \in A \text{ and } (v, u) \in A)$ but not parallel arcs (A is not a multiset).

- **Definition.** Let $D = (V, A)$ be a digraph. Then, \overline{D} denotes the **complement** of D ; this is the digraph $(V, (V \times V) \setminus A)$. Its arcs are the **non-arcs** of D .

The complement of a digraph

- **Definition.** Let $D = (V, A)$ be a digraph. Then, \bar{D} denotes the **complement** of D ; this is the digraph $(V, (V \times V) \setminus A)$. Its arcs are the **non-arcs** of D .
- **Example.**




- **Definition.** A digraph $D = (V, A)$ is **loopless** if it has no loops (i.e., no arcs (u, u)).

- **Definition.** A digraph $D = (V, A)$ is **loopless** if it has no loops (i.e., no arcs (u, u)).
- **Definition.** A loopless digraph $D = (V, A)$ is a **tournament** if it has the following property: For any distinct $u, v \in V$, exactly one of the two pairs (u, v) and (v, u) is an arc of D . In other words, a tournament is an orientation of the complete undirected graph K_V .

Tournaments

- **Definition.** A digraph $D = (V, A)$ is **loopless** if it has no loops (i.e., no arcs (u, u)).
- **Definition.** A loopless digraph $D = (V, A)$ is a **tournament** if it has the following property: For any distinct $u, v \in V$, exactly one of the two pairs (u, v) and (v, u) is an arc of D . In other words, a tournament is an orientation of the complete undirected graph K_V .
- **Examples.**



- **Definition.** Let V be a finite set. A **V -listing** will mean a list of elements of V that contains each element of V exactly once.

- **Definition.** Let V be a finite set. A **V-listing** will mean a list of elements of V that contains each element of V exactly once.
- **Definition.** Let $D = (V, A)$ be a digraph. A **Hamiltonian path** (short: **hamp**) of D means a V -listing (v_1, v_2, \dots, v_n) such that

$$(v_i, v_{i+1}) \in A \quad \text{for each } i \in \{1, 2, \dots, n-1\}.$$


In other words (for $V \neq \emptyset$), it means a path of D that contains each vertex.

- **Easy proposition (Rédei 1933):** Any tournament has a hamper.

- **Easy proposition (Rédei 1933):** Any tournament has a hamp.
- This is an easy exercise in graph theory. But Rédei proved a lot more:
- **Theorem (Rédei 1933):** Let D be a tournament. Then,
(# of hamps of D) is odd.

- **Easy proposition (Rédei 1933):** Any tournament has a hamp.
- This is an easy exercise in graph theory. But Rédei proved a lot more:
- **Theorem (Rédei 1933):** Let D be a tournament. Then,


($\#$ of hamps of D) is odd.
- **Example.** Here are some tournaments:



- Recall **Redei's Theorem**: Let D be a tournament. Then,
(# of hamps of D) is odd.
- Rédei's proof is complicated and intransparent (see Moon, *Topics on Tournaments* for an English version).

Berge's theorem

- Recall **Redei's Theorem**: Let D be a tournament. Then,
(# of hamps of D) is odd.
- Rédei's proof is complicated and intransparent (see [Moon, Topics on Tournaments](#) for an English version).
To give a more conceptual proof, Berge discovered the following:
 - Theorem (Berge 1976)**: Let D be a digraph. Then,
(# of hamps of \bar{D}) \equiv (# of hamps of D) mod 2.
 - Example.**



- Berge proves his theorem (in his *Graphs* textbook) using an elegant inclusion-exclusion argument.

Then he uses his theorem to prove Rédei's theorem via induction on the number of "inversions" (arcs directed the "wrong way").

This proof is much cleaner than Rédei's, but still far from simple.

For a detailed exposition, see [https:](https://www.cip.ifi.lmu.de/~grinberg/t/17s/5707lec7.pdf)

[//www.cip.ifi.lmu.de/~grinberg/t/17s/5707lec7.pdf](https://www.cip.ifi.lmu.de/~grinberg/t/17s/5707lec7.pdf)

.

- Berge proves his theorem (in his *Graphs* textbook) using an elegant inclusion-exclusion argument.

Then he uses his theorem to prove Rédei's theorem via induction on the number of "inversions" (arcs directed the "wrong way").

This proof is much cleaner than Rédei's, but still far from simple.

For a detailed exposition, see [https:](https://www.cip.ifi.lmu.de/~grinberg/t/17s/5707lec7.pdf)

[//www.cip.ifi.lmu.de/~grinberg/t/17s/5707lec7.pdf](https://www.cip.ifi.lmu.de/~grinberg/t/17s/5707lec7.pdf)

- **Remark.** Can we improve on Rédei's theorem even further? [MathOverflow question #232751](#) asks for the possible values of (# of hamps of D) for a tournament D . Among the numbers between 1 and 80555, the answer is "all odd numbers except for 7 and 21" (proved by bof and Gordon Royle).
Question: Are these the only exceptions?

- Independently, **Chow** (*The Path-Cycle Symmetric Function of a Digraph*, 1996) introduced a symmetric function assigned to each digraph D .
(This was inspired by Chung/Graham's cover polynomial in rook theory.)
- We only discuss a coarsening of his construction (Chow has two families of variables, and we set the second family to 0).
Question: Which of the results below can be generalized to the full version?

- **Definition.** Let $n \in \mathbb{N}$, and let I be a subset of $\{1, 2, \dots, n-1\}$. Then, we define the power series

$$L_{I,n} := \sum_{\substack{i_1 \leq i_2 \leq \dots \leq i_n; \\ i_p < i_{p+1} \text{ for each } p \in I}} x_{i_1} x_{i_2} \cdots x_{i_n} \in \mathbb{Z}[[x_1, x_2, x_3, \dots]]$$

(where the indices i_1, i_2, \dots, i_n range over $\{1, 2, 3, \dots\}$).

Remark: This is a formal power series (but becomes a polynomial if you drop all but finitely many variables). It is known as a **(Gessel's) fundamental quasisymmetric function**.

The Rédei–Berge symmetric function: definition

- **Definition.** Let $n \in \mathbb{N}$, and let I be a subset of $\{1, 2, \dots, n-1\}$. Then, we define the power series

$$L_{I,n} := \sum_{\substack{i_1 \leq i_2 \leq \dots \leq i_n; \\ i_p < i_{p+1} \text{ for each } p \in I}} x_{i_1} x_{i_2} \cdots x_{i_n} \in \mathbb{Z}[[x_1, x_2, x_3, \dots]]$$

(where the indices i_1, i_2, \dots, i_n range over $\{1, 2, 3, \dots\}$).

Remark: This is a formal power series (but becomes a polynomial if you drop all but finitely many variables).

It is known as a **(Gessel's) fundamental quasisymmetric function**.

- For instance,

$$L_{\{1\},3} = \sum_{i_1 < i_2 \leq i_3} x_{i_1} x_{i_2} x_{i_3};$$

$$L_{\{1\},4} = \sum_{i_1 < i_2 \leq i_3 \leq i_4} x_{i_1} x_{i_2} x_{i_3} x_{i_4}.$$

The Rédei–Berge symmetric function: definition

- **Definition.** Let $n \in \mathbb{N}$, and let I be a subset of $\{1, 2, \dots, n-1\}$. Then, we define the power series

$$L_{I,n} := \sum_{\substack{i_1 \leq i_2 \leq \dots \leq i_n; \\ i_p < i_{p+1} \text{ for each } p \in I}} x_{i_1} x_{i_2} \cdots x_{i_n} \in \mathbb{Z}[[x_1, x_2, x_3, \dots]]$$

(where the indices i_1, i_2, \dots, i_n range over $\{1, 2, 3, \dots\}$).

- **Definition.** Let $n \in \mathbb{N}$. Let $D = (V, A)$ be a digraph with n vertices. We define the **Redei–Berge symmetric function**


$$U_D := \sum_{w \text{ is a } V\text{-listing}} L_{\text{Des}(w,D)}, \quad n \in \mathbb{Z}[[x_1, x_2, x_3, \dots]],$$

where

$$\text{Des}(w, D) := \{i \in \{1, 2, \dots, n-1\} \mid (w_i, w_{i+1}) \in A\}$$

for each V -listing $w = (w_1, w_2, \dots, w_n)$.

- **Example:** Let



Then,

$$\begin{aligned}
 U_D &= \sum_{w \text{ is a } V\text{-listing}} L_{\text{Des}(w,D), 3} \\
 &= L_{\text{Des}((1,2,3),D), 3} + L_{\text{Des}((1,3,2),D), 3} + L_{\text{Des}((2,1,3),D), 3} \\
 &\quad + L_{\text{Des}((2,3,1),D), 3} + L_{\text{Des}((3,1,2),D), 3} + L_{\text{Des}((3,2,1),D), 3} \\
 &= L_{\{1\}, 3} + L_{\emptyset, 3} + L_{\emptyset, 3} + L_{\emptyset, 3} + L_{\{2\}, 3} + L_{\emptyset, 3} \\
 &= 4 \cdot L_{\emptyset, 3} + L_{\{1\}, 3} + L_{\{2\}, 3} \\
 &= 4 \cdot \sum_{i_1 \leq i_2 \leq i_3} x_{i_1} x_{i_2} x_{i_3} + \sum_{i_1 < i_2 \leq i_3} x_{i_1} x_{i_2} x_{i_3} + \sum_{i_1 \leq i_2 < i_3} x_{i_1} x_{i_2} x_{i_3}.
 \end{aligned}$$

The Rédei–Berge symmetric function: restatement

- We can restate the definition of U_D directly as follows:
- **Proposition.** Let $D = (V, A)$ be a digraph. Then,

$$U_D = \sum_{f: V \rightarrow \{1,2,3,\dots\}} a_{D,f} \prod_{v \in V} x_{f(v)},$$

where $a_{D,f}$ is the # of all V -listings $w = (w_1, w_2, \dots, w_n)$ such that

- we have $f(w_1) \leq f(w_2) \leq \dots \leq f(w_n)$;
- we have $f(w_i) < f(w_{i+1})$ if $(w_i, w_{i+1}) \in A$.

The Rédei–Berge symmetric function: restatement

- We can restate the definition of U_D directly as follows:
- **Proposition.** Let $D = (V, A)$ be a digraph. Then,

$$U_D = \sum_{f:V \rightarrow \{1,2,3,\dots\}} a_{D,f} \prod_{v \in V} x_{f(v)},$$

where $a_{D,f}$ is the # of all V -listings $w = (w_1, w_2, \dots, w_n)$ such that

- we have $f(w_1) \leq f(w_2) \leq \dots \leq f(w_n)$;
 - we have $f(w_i) < f(w_{i+1})$ if $(w_i, w_{i+1}) \in A$.
- This is similar (though not directly related) to P -partition enumerators and chromatic symmetric functions.

The Rédei–Berge symmetric function: restatement

- We can restate the definition of U_D directly as follows:
- **Proposition.** Let $D = (V, A)$ be a digraph. Then,

$$U_D = \sum_{f: V \rightarrow \{1,2,3,\dots\}} a_{D,f} \prod_{v \in V} x_{f(v)},$$

where $a_{D,f}$ is the # of all V -listings $w = (w_1, w_2, \dots, w_n)$ such that

- we have $f(w_1) \leq f(w_2) \leq \dots \leq f(w_n)$;
 - we have $f(w_i) < f(w_{i+1})$ if $(w_i, w_{i+1}) \in A$.
- **Remark.** We can restate the definition of $a_{D,f}$ in nicer terms. Namely, fix a digraph $D = (V, A)$ and a map $f: V \rightarrow \{1, 2, 3, \dots\}$. For any $j \in f(V)$, let $\overline{D_j}$ denote the induced subdigraph of the complement \overline{D} on the vertex set $f^{-1}(j) = \{v \in V \mid f(v) = j\}$. Then,

$$a_{D,f} = \prod_{j \in f(V)} (\# \text{ of hamps of } \overline{D_j}).$$

- Note that U_D is $\Xi_{\overline{D}}(x, 0)$ in the notations of Chow's 1996 paper.

- Note that U_D is $\Xi_{\overline{D}}(x, 0)$ in the notations of Chow's 1996 paper.
- What is U_D good for? Counting hamps (= Hamiltonian paths), for one:
- **Proposition.** Let D be a digraph. Then,

$$U_D(1, 0, 0, 0, \dots) = (\# \text{ of hamps of } \overline{D}).$$

- Note that U_D is $\Xi_{\overline{D}}(x, 0)$ in the notations of Chow's 1996 paper.
- What is U_D good for? Counting hamps (= Hamiltonian paths), for one:
- **Proposition.** Let D be a digraph. Then,

$$U_D(1, 0, 0, 0, \dots) = (\# \text{ of hamps of } \overline{D}).$$

- Thus, any results about U_D might give us information about the # of hamps!

- Note that U_D is $\Xi_{\overline{D}}(x, 0)$ in the notations of Chow's 1996 paper.
- What is U_D good for? Counting hamps (= Hamiltonian paths), for one:
- **Proposition.** Let D be a digraph. Then,

$$U_D(1, 0, 0, 0, \dots) = (\# \text{ of hamps of } \overline{D}).$$

- Thus, any results about U_D might give us information about the # of hamps!
- Formulas for U_D in some specific cases (D acyclic, D poset, D path) can be found in **Additional Problem 120 to Chapter 7 of Stanley's EC2**. Most prominently, if D is the “greater-than digraph” of a poset P , then $U_{\overline{D}}$ is the chromatic symmetric function of the incomparability graph of P .

- I called U_D the “Rédei–Berge symmetric function”, but is it actually symmetric? Yes, and in fact something better holds:
- **Definition.** For each $k \geq 1$, let

$$p_k := x_1^k + x_2^k + x_3^k + \dots$$

be the k -th **power-sum symmetric function**.

- **Theorem.** For any digraph D , we have

$$U_D \in \mathbb{Z}[p_1, p_2, p_3, \dots].$$

That is, U_D can be written as a polynomial in p_1, p_2, p_3, \dots over \mathbb{Z} .

- I called U_D the “Rédei–Berge symmetric function”, but is it actually symmetric? Yes, and in fact something better holds:
- **Definition.** For each $k \geq 1$, let

$$p_k := x_1^k + x_2^k + x_3^k + \cdots$$

be the k -th **power-sum symmetric function**.

- **Theorem.** For any digraph D , we have

$$U_D \in \mathbb{Z}[p_1, p_2, p_3, \dots].$$

That is, U_D can be written as a polynomial in p_1, p_2, p_3, \dots over \mathbb{Z} .

- Which polynomial, though?

- **Definition.** Fix a digraph $D = (V, A)$.
Let \mathfrak{S}_V be the symmetric group on the set V .
For any $\sigma \in \mathfrak{S}_V$, we let

$$p_{\text{type } \sigma} := \prod_{\gamma \text{ is a cycle of } \sigma} p_{\text{length of } \gamma}.$$

In other words, if σ has cycles of lengths a, b, \dots, k (including 1-cycles), then $p_{\text{type } \sigma} = p_a p_b \cdots p_k$.

- **Definition.** Fix a digraph $D = (V, A)$.
Let \mathfrak{S}_V be the symmetric group on the set V .
For any $\sigma \in \mathfrak{S}_V$, we let

$$p_{\text{type } \sigma} := \prod_{\gamma \text{ is a cycle of } \sigma} p_{\text{length of } \gamma}.$$

We say that a cycle γ of σ is a **D -cycle** if all the pairs $(i, \sigma(i))$ for $i \in \gamma$ are arcs of D .

- **Definition.** Fix a digraph $D = (V, A)$.
Let \mathfrak{S}_V be the symmetric group on the set V .
For any $\sigma \in \mathfrak{S}_V$, we let

$$p_{\text{type } \sigma} := \prod_{\gamma \text{ is a cycle of } \sigma} p_{\text{length of } \gamma}.$$

We say that a cycle γ of σ is a **D -cycle** if all the pairs $(i, \sigma(i))$ for $i \in \gamma$ are arcs of D .

- **Main Theorem I.** Let $D = (V, A)$ be a digraph. Set

$$\varphi(\sigma) := \sum_{\substack{\gamma \text{ is a cycle of } \sigma; \\ \gamma \text{ is a } D\text{-cycle}}} ((\text{length of } \gamma) - 1) \quad \text{for each } \sigma \in \mathfrak{S}_V.$$

Then,

$$U_D = \sum_{\substack{\sigma \in \mathfrak{S}_V; \\ \text{each cycle of } \sigma \text{ is} \\ \text{a } D\text{-cycle or a } \overline{D}\text{-cycle}}} (-1)^{\varphi(\sigma)} p_{\text{type } \sigma}.$$

- **Definition.** Fix a digraph $D = (V, A)$.
Let \mathfrak{S}_V be the symmetric group on the set V .
For any $\sigma \in \mathfrak{S}_V$, we let

$$p_{\text{type } \sigma} := \prod_{\gamma \text{ is a cycle of } \sigma} p_{\text{length of } \gamma}.$$

We say that a cycle γ of σ is a **D -cycle** if all the pairs $(i, \sigma(i))$ for $i \in \gamma$ are arcs of D .

- **Main Theorem I.** Let $D = (V, A)$ be a digraph. Set


$$\varphi(\sigma) := \sum_{\substack{\gamma \text{ is a cycle of } \sigma; \\ \gamma \text{ is a } D\text{-cycle}}} ((\text{length of } \gamma) - 1) \quad \text{for each } \sigma \in \mathfrak{S}_V.$$

Then,

$$U_D = \sum_{\substack{\sigma \in \mathfrak{S}_V; \\ \text{each cycle of } \sigma \text{ is} \\ \text{a } D\text{-cycle or a } \bar{D}\text{-cycle}}} (-1)^{\varphi(\sigma)} p_{\text{type } \sigma}.$$

- This yields the $U_D \in \mathbb{Z}[p_1, p_2, p_3, \dots]$ theorem, of course.

- **Example.** Recall our favorite example:




The cycles of D are $(2)_\sim$ and $(3)_\sim$, whereas the cycles of \bar{D} are $(1)_\sim$, $(2, 3)_\sim$, $(3, 1)_\sim$ and $(1, 3, 2)_\sim$.

Thus, the $\sum_{\substack{\sigma \in \mathfrak{S}_V; \\ \text{each cycle of } \sigma \text{ is} \\ \text{a } D\text{-cycle or a } \bar{D}\text{-cycle}}}$ sum in Main Theorem I has four

addends, corresponding to (σ written in one-line notation)

$\sigma =$	[1, 2, 3]	[3, 1, 2]	[1, 3, 2]	[3, 2, 1]
$(-1)^{\varphi(\sigma)} =$	1	1	1	1
$p_{\text{type } \sigma} =$	p_1^3	p_3	$p_2 p_1$	$p_2 p_1$

- **Example.** Recall our favorite example:




The cycles of D are $(2)_\sim$ and $(3)_\sim$, whereas the cycles of \bar{D} are $(1)_\sim$, $(2, 3)_\sim$, $(3, 1)_\sim$ and $(1, 3, 2)_\sim$.

Hence, Main Theorem I yields

$$U_D = p_1^3 + p_3 + p_2 p_1 + p_2 p_1 = p_1^3 + 2p_1 p_2 + p_3.$$

- **Example.** Another example: Let



Thus, the $\sum_{\substack{\sigma \in \mathfrak{S}_V; \\ \text{each cycle of } \sigma \text{ is} \\ \text{a } D\text{-cycle or a } \bar{D}\text{-cycle}}}$ sum in Main Theorem I has three addends, with

$\sigma =$	[1, 2, 3]	[3, 1, 2]	[3, 2, 1]
$(-1)^{\varphi(\sigma)} =$	1	1	-1
$p_{\text{type } \sigma} =$	p_1^3	p_3	$p_2 p_1$

Hence, Main Theorem I yields $U_D = p_1^3 + p_3 - p_2 p_1$.

- Recall **Main Theorem I**: Let $D = (V, A)$ be a digraph. Set

$$\varphi(\sigma) := \sum_{\substack{\gamma \text{ is a cycle of } \sigma; \\ \gamma \text{ is a } D\text{-cycle}}} ((\text{length of } \gamma) - 1) \quad \text{for each } \sigma \in \mathfrak{S}_V.$$

Then,

$$U_D = \sum_{\substack{\sigma \in \mathfrak{S}_V; \\ \text{each cycle of } \sigma \text{ is} \\ \text{a } D\text{-cycle or a } \bar{D}\text{-cycle}}} (-1)^{\varphi(\sigma)} p_{\text{type } \sigma}.$$

- Recall **Main Theorem I**: Let $D = (V, A)$ be a digraph. Set

$$\varphi(\sigma) := \sum_{\substack{\gamma \text{ is a cycle of } \sigma; \\ \gamma \text{ is a } D\text{-cycle}}} ((\text{length of } \gamma) - 1) \quad \text{for each } \sigma \in \mathfrak{S}_V.$$

Then,

$$U_D = \sum_{\substack{\sigma \in \mathfrak{S}_V; \\ \text{each cycle of } \sigma \text{ is} \\ \text{a } D\text{-cycle or a } \overline{D}\text{-cycle}}} (-1)^{\varphi(\sigma)} p_{\text{type } \sigma}.$$

- Main Theorem I yields Berge's theorem, since the sum for D and the sum for \overline{D} range over the same σ 's, and the addends only differ in sign.

- Recall **Main Theorem I**: Let $D = (V, A)$ be a digraph. Set

$$\varphi(\sigma) := \sum_{\substack{\gamma \text{ is a cycle of } \sigma; \\ \gamma \text{ is a } D\text{-cycle}}} ((\text{length of } \gamma) - 1) \quad \text{for each } \sigma \in \mathfrak{S}_V.$$

Then,

$$U_D = \sum_{\substack{\sigma \in \mathfrak{S}_V; \\ \text{each cycle of } \sigma \text{ is} \\ \text{a } D\text{-cycle or a } \bar{D}\text{-cycle}}} (-1)^{\varphi(\sigma)} p_{\text{type } \sigma}.$$

- Main Theorem I yields Berge's theorem, since the sum for D and the sum for \bar{D} range over the same σ 's, and the addends only differ in sign.
- Corollary.** Let $D = (V, A)$ be a digraph. Assume that every D -cycle has odd length. Then,

$$U_D = \sum_{\substack{\sigma \in \mathfrak{S}_V; \\ \text{each cycle of } \sigma \text{ is} \\ \text{a } D\text{-cycle or a } \bar{D}\text{-cycle}}} p_{\text{type } \sigma} \in \mathbb{N}[p_1, p_2, p_3, \dots].$$

- Main Theorem II.** Let $D = (V, A)$ be a tournament. For each $\sigma \in \mathfrak{S}_V$, let $\psi(\sigma)$ denote the number of nontrivial cycles of σ . (A cycle is called **nontrivial** if it has length > 1 .) Then,

$$U_D = \sum_{\substack{\sigma \in \mathfrak{S}_V; \\ \text{each cycle of } \sigma \text{ is a } D\text{-cycle;} \\ \text{all cycles of } \sigma \text{ have odd length}}} 2^{\psi(\sigma)} p_{\text{type } \sigma}$$

$$\in \mathbb{N}[p_1, 2p_3, 2p_5, 2p_7, \dots] = \mathbb{N}[p_1, 2p_i \mid i > 1 \text{ is odd}].$$

- **Main Theorem II.** Let $D = (V, A)$ be a tournament. For each $\sigma \in \mathfrak{S}_V$, let $\psi(\sigma)$ denote the number of nontrivial cycles of σ . (A cycle is called **nontrivial** if it has length > 1 .) Then,

$$U_D = \sum_{\substack{\sigma \in \mathfrak{S}_V; \\ \text{each cycle of } \sigma \text{ is a } D\text{-cycle;} \\ \text{all cycles of } \sigma \text{ have odd length}}} 2^{\psi(\sigma)} p_{\text{type } \sigma}$$
$$\in \mathbb{N}[p_1, 2p_3, 2p_5, 2p_7, \dots] = \mathbb{N}[p_1, 2p_i \mid i > 1 \text{ is odd}].$$

- Main Theorem II easily yields Rédei's theorem, as the only addend with $2^{\psi(\sigma)}$ odd is the $\sigma = \text{id}$ addend.

- The above corollary from Main Theorem I yields that U_D is p -positive when D has no even-length cycles. But this holds even more generally:

- The above corollary from Main Theorem I yields that U_D is p -positive when D has no even-length cycles. But this holds even more generally:
- **Main Theorem III.** Let $D = (V, A)$ be a digraph that has no cycles of length 2. Then,

$$U_D = \sum_{\substack{\sigma \in \mathcal{G}_V; \\ \text{each cycle of } \sigma \text{ is} \\ \text{a } D\text{-cycle or a } \bar{D}\text{-cycle;} \\ \text{no even-length cycle of } \sigma \text{ is} \\ \text{a } D\text{-cycle or a reversed } D\text{-cycle}}} p_{\text{type } \sigma}.$$

- The above corollary from Main Theorem I yields that U_D is p -positive when D has no even-length cycles. But this holds even more generally:
- **Main Theorem III.** Let $D = (V, A)$ be a digraph that has no cycles of length 2. Then,

$$U_D = \sum_{\substack{\sigma \in \mathcal{G}_V; \\ \text{each cycle of } \sigma \text{ is} \\ \text{a } D\text{-cycle or a } \bar{D}\text{-cycle;} \\ \text{no even-length cycle of } \sigma \text{ is} \\ \text{a } D\text{-cycle or a reversed } D\text{-cycle}}} p_{\text{type } \sigma}.$$

- **Remark.** Not all p -positive U_D 's are explained by this theorem.

- The proof of Main Theorem I is long and intricate. It might be simplifiable. Here are the main ideas.

- The proof of Main Theorem I is long and intricate. It might be simplifiable. Here are the main ideas.
- **Pólya-style lemma.** Let V be a finite set. Let $\sigma \in \mathfrak{S}_V$ be a permutation of V . Then,

$$\sum_{\substack{f: V \rightarrow \{1,2,3,\dots\}; \\ f \circ \sigma = f}} \prod_{v \in V} x_{f(v)} = p_{\text{type } \sigma}.$$

Proof. Easy exercise.

- The proof of Main Theorem I is long and intricate. It might be simplifiable. Here are the main ideas.
- **Pólya-style lemma.** Let V be a finite set. Let $\sigma \in \mathfrak{S}_V$ be a permutation of V . Then,

$$\sum_{\substack{f: V \rightarrow \{1,2,3,\dots\}; \\ f \circ \sigma = f}} \prod_{v \in V} x_{f(v)} = p_{\text{type } \sigma}.$$

- Using this lemma (and the above formula for $a_{D,f}$), we can easily reduce Main Theorem I to the following lemma:
- **Main combinatorial lemma.** Let $D = (V, A)$ be a digraph with n vertices. Let $f : V \rightarrow \{1, 2, 3, \dots\}$ be any map. Then,

$$\prod_{j \in f(V)} (\# \text{ of hamps of } \overline{D}_j) = \sum_{\substack{\sigma \in \mathfrak{S}_V; \\ \text{each cycle of } \sigma \text{ is} \\ \text{a } D\text{-cycle or a } \overline{D}\text{-cycle}; \\ f \circ \sigma = f}} (-1)^{\varphi(\sigma)},$$

where \overline{D}_j is the induced subdigraph of \overline{D} on the vertex set $f^{-1}(j)$.

- So we need to prove the **Main combinatorial lemma**: Let $D = (V, A)$ be a digraph with n vertices. Let $f : V \rightarrow \{1, 2, 3, \dots\}$ be any map. Then,

$$\prod_{j \in f(V)} (\# \text{ of hamps of } \overline{D}_j) = \sum_{\substack{\sigma \in \mathfrak{S}_V; \\ \text{each cycle of } \sigma \text{ is} \\ \text{a } D\text{-cycle or a } \overline{D}\text{-cycle}; \\ f \circ \sigma = f}} (-1)^{\varphi(\sigma)},$$

where \overline{D}_j is the induced subdigraph of \overline{D} on the vertex set $f^{-1}(j)$.

- So we need to prove the **Main combinatorial lemma**: Let $D = (V, A)$ be a digraph with n vertices. Let $f : V \rightarrow \{1, 2, 3, \dots\}$ be any map. Then,

$$\prod_{j \in f(V)} (\# \text{ of hamps of } \overline{D}_j) = \sum_{\substack{\sigma \in \mathfrak{S}_V; \\ \text{each cycle of } \sigma \text{ is} \\ \text{a } D\text{-cycle or a } \overline{D}\text{-cycle}; \\ f \circ \sigma = f}} (-1)^{\varphi(\sigma)},$$

where \overline{D}_j is the induced subdigraph of \overline{D} on the vertex set $f^{-1}(j)$.

- Work on each “level set” $f^{-1}(j)$ separately:

Main combinatorial lemma (simplified). Let $D = (V, A)$ be a digraph. Then,

$$(\# \text{ of hamps of } \overline{D}) = \sum_{\substack{\sigma \in \mathfrak{S}_V; \\ \text{each cycle of } \sigma \text{ is} \\ \text{a } D\text{-cycle or a } \overline{D}\text{-cycle}}} (-1)^{\varphi(\sigma)}.$$

- So we need to prove the **Main combinatorial lemma**: Let $D = (V, A)$ be a digraph with n vertices. Let $f : V \rightarrow \{1, 2, 3, \dots\}$ be any map. Then,

$$\prod_{j \in f(V)} (\# \text{ of hamps of } \overline{D}_j) = \sum_{\substack{\sigma \in \mathfrak{S}_V; \\ \text{each cycle of } \sigma \text{ is} \\ \text{a } D\text{-cycle or a } \overline{D}\text{-cycle}; \\ f \circ \sigma = f}} (-1)^{\varphi(\sigma)},$$

where \overline{D}_j is the induced subdigraph of \overline{D} on the vertex set $f^{-1}(j)$.

- Work on each “level set” $f^{-1}(j)$ separately:
Main combinatorial lemma (simplified). Let $D = (V, A)$ be a digraph. Then,

$$(\# \text{ of hamps of } \overline{D}) = \sum_{\substack{\sigma \in \mathfrak{S}_V; \\ \text{each cycle of } \sigma \text{ is} \\ \text{a } D\text{-cycle or a } \overline{D}\text{-cycle}}} (-1)^{\varphi(\sigma)}.$$

- This can be proved using a nontrivial exclusion-inclusion.

- To prove Main Theorems II and III, start with the Main Theorem I sum, and combine σ 's into equivalence classes by reversing certain cycles:
 - For Main Theorem II, call two permutations in \mathfrak{S}_V equivalent if one can be obtained from the other by reversing (nontrivial) cycles. This turns D -cycles into \overline{D} -cycles and vice versa. The equivalence class of σ has $2^{\psi(\sigma)}$ elements if σ has no 2-cycles. Their addends in the sum cancel out if σ has an even-length cycle; otherwise they are all equal and sum up to $2^{\psi(\sigma)} p_{\text{type } \sigma}$.

- To prove Main Theorems II and III, start with the Main Theorem I sum, and combine σ 's into equivalence classes by reversing certain cycles:
 - For Main Theorem II, call two permutations in \mathfrak{S}_V equivalent if one can be obtained from the other by reversing (nontrivial) cycles. This turns D -cycles into \overline{D} -cycles and vice versa. The equivalence class of σ has $2^{\psi(\sigma)}$ elements if σ has no 2-cycles. Their addends in the sum cancel out if σ has an even-length cycle; otherwise they are all equal and sum up to $2^{\psi(\sigma)} p_{\text{type } \sigma}$.
 - For Main Theorem III, call a necklace $(v_1, v_2, \dots, v_k)_{\sim}$ **risky** if its length k is even and either it or its inverse is a D -cycle. Call two permutations in \mathfrak{S}_V equivalent if one can be obtained from the other by reversing risky cycles. The equivalence class of σ has $2^{r(\sigma)}$ elements, where $r(\sigma)$ is the number of risky cycles of σ . Their addends in the sum cancel out if σ has a risky cycle; otherwise there is only one of them.

- The proof of Main Theorem I is detailed in the preprint (<https://arxiv.org/abs/2307.05569>); the proofs of II and III are outlined.

These would make a good project for formalization (Coq, Lean, etc.): only elementary combinatorics but some tricky reasoning with cycles and sums.

- Rédei's theorem determines the # of hamps of a tournament D modulo 2. What about $\text{mod } 4$?

- Rédei's theorem determines the # of hamps of a tournament D modulo 2. What about mod 4?
- **Theorem.** Let D be a tournament. Then,

$$\begin{aligned} & (\# \text{ of hamps of } D) \\ & \equiv 1 + 2(\# \text{ of nontrivial odd-length } D\text{-cycles}) \pmod{4}. \end{aligned}$$

Here, “nontrivial” means “having length > 1 ”.

- We can prove this using Main Theorem II. We have not seen this anywhere in the literature.

Main Theorem I “generalized”

- Main Theorem I can be rewritten without speaking about digraphs:

Main Theorem I “generalized”

- **Theorem.** Let $n \in \mathbb{N}$, and let V be an n -element set. Let \mathbf{k} be a commutative ring.

For any $a = (i, j) \in V \times V$, we fix an element $t_a = t_{(i,j)} \in \mathbf{k}$ and set $s_a := t_a + 1$.

We define the **deformed Redei–Berge symmetric function**

$$\tilde{U}_t := \sum_{\substack{w=(w_1, w_2, \dots, w_n) \\ \text{is a } V\text{-listing}}} \sum_{i_1 \leq i_2 \leq \dots \leq i_n} \left(\prod_{\substack{k \in [n-1]; \\ i_k = i_{k+1}}} s_{(w_k, w_{k+1})} \right) x_{i_1} x_{i_2} \cdots x_{i_n} \\ \in \mathbf{k} [[x_1, x_2, x_3, \dots]].$$

Then,

$$\tilde{U}_t = \sum_{\sigma \in \mathfrak{S}_V} \left(\prod_{\gamma \text{ is a cycle of } \sigma} \left(\prod_{i \in \gamma} s_{(i, \sigma(i))} - \prod_{i \in \gamma} t_{(i, \sigma(i))} \right) \right) p_{\text{type } \sigma}.$$

Main Theorem I “generalized”

- **Theorem.** Let $n \in \mathbb{N}$, and let V be an n -element set. Let \mathbf{k} be a commutative ring.

For any $a = (i, j) \in V \times V$, we fix an element $t_a = t_{(i,j)} \in \mathbf{k}$ and set $s_a := t_a + 1$.

We define the **deformed Redei–Berge symmetric function**

$$\tilde{U}_t := \sum_{\substack{w=(w_1, w_2, \dots, w_n) \\ \text{is a } V\text{-listing}}} \sum_{i_1 \leq i_2 \leq \dots \leq i_n} \left(\prod_{\substack{k \in [n-1]; \\ i_k = i_{k+1}}} s_{(w_k, w_{k+1})} \right) x_{i_1} x_{i_2} \cdots x_{i_n} \\ \in \mathbf{k} [[x_1, x_2, x_3, \dots]].$$

Then,

$$\tilde{U}_t = \sum_{\sigma \in \mathfrak{S}_V} \left(\prod_{\gamma \text{ is a cycle of } \sigma} \left(\prod_{i \in \gamma} s_{(i, \sigma(i))} - \prod_{i \in \gamma} t_{(i, \sigma(i))} \right) \right) p_{\text{type } \sigma}.$$

- This generalizes Main Theorem I (set each t_a to 0 or -1), but also follows from it by multilinearity.

- **Richard P. Stanley** for the obvious reasons.
- **Hsin Chieh Liao, Anna Pun, Bruce Sagan, Mike Zabrocki** for helpful comments.
- **the organizers** for the invitation.
- **you** for your patience.