

Sweedler's duals and Schützenberger's calculus

G erard H. E. DUCHAMP AND CHRISTOPHE TOLLU.*

Abstract

We describe the problem of Sweedler's duals for bialgebras as essentially characterizing the domain of the transpose of the multiplication. This domain is the set of what could be called "representative linear forms" which are the elements of the algebraic dual which are also representative on the multiplicative semigroup of the algebra.

When the algebra is free, this notion is indeed equivalent to that of rational functions of automata theory. For the sake of applications, the range of coefficients has been considerably broadened, i.e. extended to semirings, so that the results could be specialized to the boolean and multiplicity cases. This requires some caution (use of "positive formulas", iteration replacing inversion, rank replacing dimension for instance). For the theory and its applications has been created a rational calculus which can, in return, be applied to harness Sweedler's duals. A new theorem of rational closure and application to Hopf algebras of use in Physics and Combinatorics is provided. ⁰

1 Representative functions on a semigroup

The aim of this paragraph is to discuss the dualization of bi-algebras and Hopf algebras. This problem, solved by Sweedler's duals, is the following.

Let $(B, \cdot, \Delta, 1_B, \varepsilon)$ be a k -bialgebra ; we know that, if B is finite-dimensional (resp. graded in finite dimensions), the dual (resp. graded dual) endowed with the transpose structure is a bialgebra and that, in case B is a Hopf algebra this statement carries over. Now the question can be asked.

What is the good notion of retracted dual for the general (i.e. ungraded finite or infinite dimensional) case ?

Analysing the dualization of the structural operations $(\cdot, \Delta, 1_B, \varepsilon)$ of B , one sees at once that only the dualization of the multiplication is problematic as, in the general case, the codomain of the transpose of \cdot is larger than $B^* \otimes B^*$.

The first result follows (and somehow extends) [1]. To state it, we need the notion of (left and right) shifts of functions on a semigroup. Let k be a field, (S, \cdot) a semigroup and $f \in k^S$. For each $s \in S$ define $f_s : x \rightarrow f(sx)$ (right shift of f) and ${}_s f : x \rightarrow f(xs)$ (left shift of f), ${}_s f_t : x \rightarrow f(txs)$ (bi-shift of f). Then, we have.

*LIPN - UMR 7030 CNRS - Universit  Paris 13 F-93430 Villetaneuse, France

⁰Version du 03.05.07 compil e le 29-04-2008 17:12

Proposition 1.1 (see also [1] paragraph 2.2) Let k be a field, (S, \cdot) a semigroup and $f \in k^S$. The following are equivalent :

- i) The family $(f_s)_{s \in S}$ is of finite rank in k^S
- ii) The family $({}_s f)_{s \in S}$ is of finite rank in k^S
- iii) The family $({}_s f_t)_{s, t \in S}$ is of finite rank in k^S
- iv) There exist a double (finite) family $(g_i, h_i)_{1 \leq i \leq n}$ of functions such that

$$\left(\forall x, y \in S \right) \left(f(xy) = \sum_{i=1}^n g_i(x) h_i(y) \right) \quad (1)$$

- v) There exists $\lambda \in k^{1 \times n}, \gamma \in k^{n \times 1}$ and $\mu : (S, \cdot) \mapsto (k^{n \times n}, \times)$ a morphism of semigroups such that $(\forall s \in S)(f(s) = \lambda \mu(s) \gamma)$.

Moreover, if S admits a neutral (i.e. is a monoid), μ of (v) above can be chosen to be the unity matrix.

Proof — Omitted

The elements of k^S which fulfill the above conditions will be called *representative functions* on S and denoted $R(k, S, \cdot)$.

Remark 1.2 i) When k is only a PID, proposition (1.1) above still holds with the five equivalent conditions, orbits and ranks being computed in \bar{k}^S (\bar{k} is the fraction field of k).
 ii) If S is finite, $R(k, S, \cdot) = k^S$ and if S is a group, one has

$$R(k, S, \cdot) = k^S \iff S \text{ is finite}$$

iii) If S is a semigroup, the equivalences above are false in general as shown by the following counterexample. Let G be a finite group and endow $S = \mathbb{N} \times G$ with the law $(n, g) * (m, h) = (0, gh)$. It can be easily checked that $(S, *)$ is a semigroup and that $R(k, S, \cdot) = k^S$.

iv) When W is a shift-invariant space and $f \in W \cap R(k, S, \cdot)$, the families $(f_s)_{s \in S}, ({}_s f)_{s \in S}, ({}_s f_t)_{s, t \in S}$ are of course in $W \cap R(k, S, \cdot)$ (and are of finite rank). Two useful examples of such “relative representative functional spaces” are with $W = C(S)$ (continuous functions) and $W = S^*$ (linear forms, S being an algebra).

v) If $T \subset S$ is a subsemigroup with finite set-theoretical complement ($S \setminus T$ is finite) then $f \in k^S$ is representative iff $f|_T$ is so. In particular, if $S = T^{(1)}$, the monoid obtained by adjunction on a unity to T , one has that $f \in k^S$ is representative iff $f|_T$ is.

2 Semirings

Throughout the text “monoid” stands for “semigroup with unit”.

Semirings are the structures adapted to matrix (with unity) computation. A semiring $(k, +, \times)$ consists of the following data

- a set k
- two binary laws $+, \times$ on k

such that

- $(k, +)$ and (k, \times) are monoids, the first being commutative, their neutrals will be denoted respectively 0_k and 1_k
- \times is left and right distributive over $+$
- 0_k is an annihilator i. e. $(\forall x \in k)(0_k \cdot x = x \cdot 0_k = 0_k)$

Example 2.1 *i) Any ring.*

ii) The boolean semiring $\mathbb{B} = \{0, 1\}$ endowed with the laws $x \oplus y = x + y - xy$ and $x \otimes y = xy$.

iii) The semiring $([-\infty, +\infty[, \max, +)$, called in the literature “(max,plus)-semiring”.

iv) In the semiring $([0, +\infty[, +, \times)$, the laws are continuous at infinity and then can be completed. We obtain a semiring $([0, +\infty], +, \times)$ which is suited for multiplicities arising in repeated additions of positive values during iterations.

Example 2.2 *The following example is fundamental and will be used in the definition of CM-modules. Let $(M, +)$ be a commutative monoid, then $(\text{End}(M), +, \circ)$ (defined as for the case when M is a group) is a semiring. The units being respectively, as in the group case, the constant mapping $M \mapsto \{0_M\}$ for $+$ and Id_M for \circ .*

The structure of semiring defines a category larger than that of rings, the morphisms being defined similarly. Let $(k_i, +_i, \times_i)$, $i = 1, 2$ be two semirings, a mapping $\phi : k_1 \mapsto k_2$ is called a morphism of semirings iff it is a morphism for the two structures of monoids (additive and multiplicative), then compatible with the laws and units of k_1 and k_2 .

The definition of modules (here called CM-modules as they are constructed on Commutative Monoids as vector structure) follows also the classical pattern.

The structure of a (left) k -CM-module is given by the following data

- a commutative monoid $(M, +)$
- a morphism (the scaling morphism) of semirings $s : k \mapsto \text{End}(M)$.

The structure of (right) k -CM-module is defined by replacing $\text{End}(M)$ by $\text{End}^{op}(M)$ the opposite semiring (constructed with the opposite multiplicative law). Bi- and multimodules are defined as in [3] and follow the general philosophy of “structures with operators”.

Example 2.3 *Let X be a set, then k^X , the set of all functions $X \mapsto k$ is naturally endowed with a structure of $k - k$ bimodule defined as in the case when k is a ring. So is $k^{(X)}$, the set of finitely supported functions of k^S (actually a sub- k - k bimodule of k^S).*

The free monoid generated by a set X (finite or infinite) is the set of words (i.e. finite sequences of elements of X comprising the empty one denoted by 1_{X^*}) endowed with the concatenation law.

3 Shift operators and rational closure

Let (M, \cdot) be a (commutative or not) monoid. For a function $f : M \mapsto k$ and $a \in M$, we define the following shift operators ([Reu, Eil, Abe])

- $f_a : x \rightarrow f(ax)$ (right shift)
- ${}_a f : x \rightarrow f(xa)$ (left shift)

We also have to describe the analog, for CM-algebras, of *full subalgebras* and *full subalgebraic closures* (see [4] Ch. 1.1.4) and this requires the notion of *summability* [2].

Definition 3.1 *A family $\{f_i\}_{i \in I}$ of functions $M \mapsto k$ is called summable iff, for each $m \in M$, $\{f_i(m)\}_{i \in I}$ is finitely supported. Then, the mapping $m \mapsto \sum_{i \in I} f_i(m)$ is denoted $\sum_{i \in I} f_i$ and called the sum of $\{f_i\}_{i \in I}$.*

As a consequence, it is easily checked that, if M is *locally finite* ([14] Vol. A VII.4) and $f : M \mapsto k$ is without constant term (i.e. $f(1_M) = 0_k$), then the family $\{f^n\}_{n \in \mathbb{N}}$ (convolutional powers) is summable and its sum

$$\sum_{n \in \mathbb{N}} f^n \tag{2}$$

will be denoted f^* and called the *star* of f .

Note 3.2 *There is a lot of literature about the star problem (see [19, 20]). For a general discussion of star-type solutions in a semiring, see [12].*

Now, we are in the position of stating the Kleene-Schützenberger theorem.

Theorem 3.3 *Let $M = X^*$ be a free monoid, k a semiring and $f \in k^M$. The following are equivalent*

- i) the family $(f_w)_{w \in M}$ belongs to a finitely generated shift-invariant left-submodule*
- ii) the family $({}_w f)_{w \in M}$ belongs to a finitely generated shift-invariant right-submodule*
- iii) there exist a row $\lambda \in k^{1 \times n}$, a column $\gamma \in k^{n \times 1}$ and a representation (of monoids) $\mu : M \mapsto (k^{n \times n}, \times)$ such that $(\forall w \in M)(f(w) = \lambda \mu(w) \gamma)$.*
- iv) if X is finite, f lies in the rational closure of X (i.e. the smallest subalgebra of $k \ll X \gg$ closed under the star operation and containing X).*

Remark 3.4 *i) One can remove the hypothesis of freeness of M if k is a field. Indeed, in this case, the submodule can be taken as generated by the shifts (right or left) of f and the representation is automatically compatible with the relations of M .*

ii) Here the star is used as the localization at one (i.e. with positive formulas) of the inverse function. Indeed, with coefficients in a ring, if we are at the neighbourhood of 1, the condition $(1 - x)(1 + y) = 1$ is equivalent to $y = x + xy$ (and $y = x + yx$). These self-reproducing positive conditions are taken as the definition of “ y is a star of x ” in a semiring (see [12]).

iii) The condition (iv) in theorem (3.3) is known under the name of Kleene-Schützenberger theorem as, when k is specialized to \mathbb{B} , this is actually Kleene’s theorem. In this sense,

this theorem lies at the frontier of harmonic analysis (the set of representative functions is dense in the Fourier space of compact groups), spectral theory (the notion of full subalgebra closure comes from this theory) and theoretical computer science (the notion of star was developed as a computational model of iteration and the notion of a semiring was developed to cope with general scalars as various as the ones arising in stochastic automata theory and the shortest path problem).

In the general case (X finite or not), Kleene-Schützenberger's theorem has to be modified as follows.

Theorem 3.5 *Let $M = X^*$ be a free monoid, k a semiring and $f \in k^M$. The following are equivalent*

- i) the family $(f_w)_{w \in M}$ belongs to a finitely generated shift-invariant left-submodule*
- ii) the family $({}_w f)_{w \in M}$ belongs to a finitely generated shift-invariant right-submodule*
- iii) there exist a row $\lambda \in k^{1 \times n}$ a column $\gamma \in k^{n \times 1}$ and a representation (of monoids) $\mu : M \mapsto (k^{n \times n}, \times)$ such that $(\forall w \in M)(f(w) = \lambda \mu(w) \gamma)$.*
- iv) the function f lies in the rational closure of $\overline{kX} = \{\sum_{x \in X} \alpha(x)x\}_{\alpha \in k^X}$ (i.e. the smallest subalgebra of $k \langle\langle X \rangle\rangle$ closed under the star operation which contains \overline{kX}).*

Remark 3.6 *The rational closure of X is, in fact, the intersection of the set of elements characterized by (i-iii) i.e. Sweedler's dual of $k\langle X \rangle$ and the algebra $\bigcup_{\substack{F \subset X \\ F \text{ finite}}} k\langle\langle F \rangle\rangle$ of the series whose support involves a finite alphabet.*

4 Rational expressions

The construction of [6] was localized at zero, we extend it here to any localization i.e. for any mapping $\Lambda : X \mapsto k$.

As the rational closure involves a unitary law (the star) partially defined, the definition of universal formulas for this closure needs some caution. Indeed, we need to build in parallel a "character" (the constant term) *const* so that all proper expressions should have a star.

One first defines, as in [6] the completely free expressions (or formulas) as the terms of the universal algebra defined on $X \cup \{0_E\}$ (0_E , which does not belong to X will serve as a null or void expression and be mapped to the zero series). This algebra will be denoted $\mathcal{E}^{cf}(X, k)$. More precisely

- If $x \in X \cup \{0_E\}$ then $x \in \mathcal{E}^{cf}(X, k)$.
- If $E, E_i \in \mathcal{E}^{cf}(X, k)$, $i = 1, 2$ and $\lambda \in k$ then

$$\begin{aligned} E_1 + E_2 &\in \mathcal{E}^{cf}(X, k), \quad E_1 \cdot E_2 \in \mathcal{E}^{cf}(X, k) \\ \lambda E &\in \mathcal{E}^{cf}(X, k), \quad E\lambda \in \mathcal{E}^{cf}(X, k). \end{aligned}$$

- If $E \in \mathcal{E}^{cf}(X, k)$ then $E^* \in \mathcal{E}^{cf}(X, k)$.

The partial function *const* : $\mathcal{E}^{cf}(X, k) \mapsto k$ (constant term) is constructed as follows from the values *const*(x) = $\Lambda(x)$.

1. If $x \in X$ $const(x) = \Lambda(x)$ and $const(0_{\mathcal{E}}) = 0_k$.

2. If $E, E_i \in \mathcal{E}^{cf}(X, k)$, $i = 1, 2$ and $\lambda \in k$ then

$$const(E_1 + E_2) = const(E_1) + const(E_2), \quad const(E_1 \cdot E_2) = const(E_1) \cdot const(E_2) \\ const(\lambda E) = \lambda const(E), \quad const(E\lambda) = const(E)\lambda.$$

3. If $const(E) = 0_k$ then $const(E^*) = 1_k$.

The domain of $const$ will be called *rational expressions* and denoted $\mathcal{E}_{\Lambda}(X, k)$. For example $0_k^* \in \mathcal{E}_{\Lambda}(X, k)$.

Let now $\Theta : X \mapsto k\langle\langle A \rangle\rangle$ be a mapping such that,

$$(\forall x \in X)(\Theta(x)[1_{A^*}] = \Lambda(x))$$

Following recursively (1-2-3) above, we can construct a polymorphism $\phi_{\Theta} : \mathcal{E}_{\Lambda}(X, k) \mapsto k\langle\langle A \rangle\rangle$ which is a morphism for the laws (2 internal and 2 external) and the star. Moreover $\delta_{1_{A^*}} \circ \phi_{\Theta} = const$ (i.e. $const$ can be considered as a “constant term function” for the expressions). The image of ϕ_{Θ} is exactly the rational closure of the set $\{\Theta(x)\}_{x \in X}$.

5 Dual laws

Let $\Delta : \mathcal{A} \mapsto \mathcal{A} \times \mathcal{A}$ be any comultiplication (i.e. \mathcal{A} is a k -coalgebra). It is known that its dual $(\mathcal{A}^*, {}^t\Delta)$ is an algebra and if \mathcal{A} is coassociative (resp. cocomutative, counital), \mathcal{A} is associative (resp. commutative, unital) [1].

We would like here to enlarge the framework of [13].

If \mathcal{A} is an algebra, let us call *dual law* on \mathcal{A}^* a law of the form ${}^t\Delta$ for some (not necessarily coassociative) comultiplication on \mathcal{A} .

In [13] were considered the *dual laws* on $k\langle\langle X \rangle\rangle \simeq k\langle X \rangle^*$ in order to prove that the Hadamard and Infiltration products, which were known to preserve rationality, were essentially the only (along with an interpolation between the two) alphabetic (associative and unital) dual laws between series. As we will see, the notion of *dual law* provides an implementation scheme for the automata so that the rationality preservation is naturally effective.

Theorem 5.1 *Let \mathcal{A} be a k -algebra and $\Delta : \mathcal{A} \mapsto \mathcal{A} \otimes \mathcal{A}$ be a comultiplication which is a morphism of algebras. Then*

i) *If k is a field, Sweedler’s dual \mathcal{A}° of \mathcal{A} is closed under the dual law ${}^t\Delta$.*

ii) *If k is a semiring and $\mathcal{A} = k\langle X \rangle$, \mathcal{A}° is closed under the dual law ${}^t\Delta$.*

Note 5.2 *i) The theorem is no longer true if $\Delta : \mathcal{A} \mapsto \mathcal{A} \otimes \mathcal{A}$ is arbitrary (i.e. not necessarily a morphism) as shows the following counterexample. With $\Delta : \mathbb{Q}[x] \mapsto \mathbb{Q}[x] \otimes \mathbb{Q}[x]$ such that $\Delta(x) = \frac{x^n}{n!} \otimes \frac{x^n}{n!}$, one has*

$${}^t\Delta\left(\frac{1}{1-x}, \frac{1}{1-x}\right) = exp(x).$$

ii) *In (i) above, the restriction on scalars (to be a field) can be extended to inductive limits of PIDs.*

iii) Other coproducts than morphisms can preserve rationality. For example, let $\Delta : k\langle X \rangle \mapsto k\langle X \rangle \otimes k\langle X \rangle$ be a morphism and $\Delta_1 : k\langle X \rangle \mapsto k\langle X \rangle \otimes k\langle X \rangle$ be a linear mapping which coincides with Δ except for a finite number of words of X^* . It can be checked that Δ_1 , although not a morphism, preserves rationality.

6 Bialgebras

Let us now return to the case of a bialgebra $(B, \cdot, \Delta, 1_B, \varepsilon)$. The following proposition says that, if a linear form on B is transformed by ${}^t m$ ($m : B \otimes B \mapsto B$ is just the multiplication mapping) into an element of $B^* \otimes B^*$, it must be of the (iv) form. For the moment, let us state this in a precise form.

Proposition 6.1 *Let $(B, m, \Delta, 1_B, \varepsilon)$ be a bialgebra and $f \in B^*$.*

1) *The following are equivalent*

i) ${}^t m(f) \in B^* \otimes B^*$ (for the canonical embedding $B^* \otimes B^* \hookrightarrow (B \otimes B)^*$)

ii) $f \in R(k, B, \cdot)$

iii) $\ker(f)$ contains a finite-codimension one-sided ideal

iv) $\ker(f)$ contains a finite-codimension two-sided ideal

v) *There exist $\lambda \in k^{1 \times n}$, $\gamma \in k^{n \times 1}$ and $\mu : (B, +, \cdot) \mapsto (k^{n \times n}, +, \times)$ a morphism of k -algebras (associative with units) such that $(\forall x \in B)(f(x) = \lambda \mu(x) \gamma)$.*

2) *Moreover, let B^0 be the set of linear forms as above, then $(B^0, {}^t \Delta, {}^t m, {}^t \varepsilon, {}^t 1_B)$ is a bialgebra and if B admits an antipode σ (i.e. is a Hopf algebra), one has ${}^t \sigma(B^0) \subset B^0$ and $(B^0, {}^t \Delta, {}^t m, {}^t \varepsilon, {}^t 1_B, {}^t \sigma)$ is a Hopf algebra.*

7 An application of rational expressions to Combinatorial Physics

In a joint work with J. Katriel, one of us gave the solution of the problem of matrix coefficients in the Fock space of carriers between two levels. We give here a Brief review of these continued fractions-type formulas and provide a sketch of their proof.

In [23] was considered, as Fock space, a general vector space V over a field k with basis $|e_n\rangle$ $n = 0, 1, \dots$, equipped with its natural grading

$$V = \bigoplus_{n \in \mathbf{Z}} V_n \text{ with } V_n := k e_n; V_{-n-1} := (0) \text{ for } n \geq 0 \quad (3)$$

and scalar product defined by $\langle e_n | e_m \rangle = \delta_{n,m}$. and f, g two linear operators in V of degrees $-1, +1$, respectively. Generically, they read

$$\text{for } n \geq 0; f|e_0\rangle := 0; f|e_{k+1}\rangle = \alpha_{k+1}|e_k\rangle; g|e_k\rangle := \beta_{k+1}|e_{k+1}\rangle \quad (4)$$

We consider the words in f, g :

$$w(f, g) := f^{p_1} g^{q_1} f^{p_2} g^{q_2} \dots f^{p_n} g^{q_n} \quad (5)$$

the degree (excess) of which is $\sum_{k=1}^n (q_k - p_k)$ (this is, for algebraists, the degree of the graded operator $f^{p_1} g^{q_1} f^{p_2} g^{q_2} \dots f^{p_n} g^{q_n}$). This provides a representation μ of a two letters

free monoid $\{b_-, b_+\}^*$ on V by $\mu(b_-) = f$; $\mu(b_+) = g$, which is graded for the weight on $\{b_-, b_+\}^*$. In order to keep the reading of a word from left to right, one performs the the action on the right. Thus V becomes a $\{b_-, b_+\}^*$ right module by

$$e_0 \cdot b_- = 0 ; e_{n+1} \cdot b_- = \alpha_{n+1} e_n ; e_n \cdot b_+ = \beta_{n+1} e_{n+1} \quad (6)$$

one is interested by the matrix elements

$$\langle e_n \cdot \{b_-, b_+\}^i | e_m \rangle = \omega_{n \rightarrow m}^{(i)} . \quad (7)$$

The following proposition gives the support i. e. the set of words w of length i such that $\langle e_n \cdot w | e_m \rangle \neq 0$. Define $W_{n \rightarrow m}^{(i)}$ as follows

$$W_{n \rightarrow m}^{(i)} = \{w \in \{b_-, b_+\}^i \mid (\pi_e(w) = m - n) \text{ and } (w = uv \implies \pi_e(u) \geq -n)\} . \quad (8)$$

The following proposition characterizes $W_{n \rightarrow m}^{(i)}$ is an universal transporter between level n and level m .

Proposition 7.1 *i) If the all the weights α_n ; $n \geq 1$, β_n , $n \geq 0$ are not zero $W_{n \rightarrow m}^{(i)}$ is exactly the set of words of length i such that $\langle e_n \cdot w | e_m \rangle \neq 0$.*

ii) In all cases the latter is a subset of $W_{n \rightarrow m}^{(i)}$ i. e.

$$\{w \in \{b_-, b_+\}^i \mid \langle e_n \cdot w | e_m \rangle \neq 0\} \subset W_{n \rightarrow m}^{(i)} \quad (9)$$

Indeed, $W_{n \rightarrow m}^{(i)}$ admits factorizations as “noncommutative continued fractions”. In order to do this, we recall for the reader the definition of Dyck, positive Dyck and negative restricted (by depth) Dyck codes respectively

$$\begin{aligned} D &= \{w \in \pi_e^{-1}(0) \mid w = uv \text{ and } \varepsilon \notin \{u, v\} \implies \pi_e(u) \neq 0\} \\ D_+ &= \{w \in D \mid w = uv \text{ and } \varepsilon \notin \{u, v\} \implies \pi_e(u) > 0\} \\ D_-^{(n)} &= \{w \in D_- \mid \min_{uv=w} (\pi_e(u)) = -n\} . \end{aligned} \quad (10)$$

Now, identifying the subsets defined previously with their characteristic series in $\mathbb{N}\langle\langle b_-, b_+ \rangle\rangle$, one has the following factorizations (here, equalities between rational expressions of b_\pm, D_+ and $D_-^{(n)}$).

Proposition 7.2 *Set $W_{n \rightarrow m} = \sum_{i \geq 0} W_{n \rightarrow m}^{(i)}$.*

1) One has the following factorizations for $W_{n \rightarrow n+k}$

$$\begin{aligned} (D_+ + D_-^{(n)})^* (b_+ D_+^*)^k &= \left[\prod_{i=0}^{\rightarrow (k-1)} ((D_-^{(n+i)})^* b_+) \right] (D_+ + D_-^{(n+k)})^* \text{ if } k \geq 0 \\ (D_+ b_-)^k (D_+ + D_-^{(n+k)})^* &= (D_+ + D_-^{(n)})^* \left[\prod_{i=1}^{\rightarrow (-k)} ((b_- (D_-^{(n-i)})^*) \right] \text{ otherwise} \end{aligned} \quad (11)$$

2) One has the following self-reproducing equations for the Dyck codes

$$D_+ = b_+ (D_+)^* b_- ; D_-^{(n)} = b_- (D_-^{(n-1)})^* b_+ ; D_-^{(0)} = \emptyset \quad (12)$$

The statements of this proposition can be considered as a “noncommutative continued fraction” expansion of $W_{n \rightarrow m}$.

As a result [23], with

$$\mu(W_{m-n}^{(i)}|n) = \omega_{n \rightarrow m}^{(i)}|m) ; T_{n \rightarrow n+k} := \sum_{i \geq 0} t^i \omega_{n \rightarrow n+k}^{(i)} , \quad (13)$$

using proposition (7.2) and the representation $\mu(b \pm) = tb \pm$, one can expand $T_{n \rightarrow n+k}$ as a product of continued fractions. Let

$$F_n^+ = \frac{1}{1 - \frac{t^2 \alpha_{n+1} \beta_{n+1}}{1 - \frac{t^2 \alpha_{n+2} \beta_{n+2}}{1 - \frac{t^2 \alpha_{n+3} \beta_{n+3}}{1 - \dots}}}} = \frac{1}{1 - E_n^+}$$

$$F_n^- = \frac{1}{1 - \frac{t^2 \alpha_n \beta_n}{1 - \frac{t^2 \alpha_{n-1} \beta_{n-1}}{1 - \frac{t^2 \alpha_{n-2} \beta_{n-2}}{1 - \dots}}}} = \frac{1}{1 - E_n^-} \quad (14)$$

and

$$F_n = \frac{1}{1 - E_n^+ - E_n^-} . \quad (15)$$

Then, if $k \geq 0$, we obtain

$$T_{n \rightarrow n+k} = t^k F_{n+k} \prod_{i=0}^{k-1} F_{n+i}^- = t^k F_n \prod_{i=1}^k F_{n+i}^+ \quad (16)$$

and, if $k \leq 0$

$$T_{n \rightarrow n+k} = t^{-k} F_{n+k} \prod_{i=0}^{-k-1} F_{n-i}^+ = t^{-k} F_n \prod_{i=1}^{-k} F_{n-i}^- \quad (17)$$

8 Conclusion

The dualization problem solved by Sweedler’s duals has striking relations with language theory (for this last point, see [2]). A true “engineer-like calculus” was developed to handle the rational closure mentioned above. This set of formulas is mainly based on a recursion to compute the “star of a matrix” (the formulas and a complete discussion can be found in [12] and are reminiscent of general formulas giving the inverse of a matrix decomposed in blocks). This calculus is powerful enough to be the main ingredient in investigating rationality properties in various domains (see [9, 11] for noncommutative geometry and [2, 14] for automata theory) and sufficiently expressive to give exact developments of some transfer coefficients in Combinatorial Physics [23]. These rational expressions are generic in the sense that any Sweedler’s dual can be described by them. In this way, like for Dirac’s notation, we can hope to inherit the computational skill developed since 40 years of practice.

References

- [1] E. ABE Hopf algebras. Cambridge (1980)
- [2] J. BERSTEL, C. REUTENAUER Rational series and their languages EATCS Monographs on Theoretical Computer Science. Springer (1988).
- [3] BOURBAKI N., *Algebra, chapter III*, Springer
- [4] BOURBAKI N., *Théories spectrales*, Hermann
- [5] A. CAYLEY, On certain results related to quaternions. Phil. Mag. **26**, 141-145 (1845).
- [6] CHAMPARNAUD J.-M., DUCHAMP G. *Derivatives of rational expressions and related theorems*, T.C.S. **313** 31 (2004).
- [7] V. CHARI, A. PRESSLEY, A guide to quantum groups. Cambridge (1994).
- [8] P.M. COHN, Free rings and their relations. Acad. Press (1985).
- [9] A. CONNES, Noncommutative geometry. Acad. Press (1994).
- [10] G. CAUCHON, Séries de Malcev-Neumann sur le groupe libre et questions de rationalité, TCS, **98**, pp79-97 (1992).
- [11] DUCHAMP G., REUTENAUER C., *Un critère de rationalité provenant de la géométrie noncommutative* Invent. Math. **128** 613-622. (1997).
- [12] GÉRARD DUCHAMP, HATEM HADJ KACEM, ERIC LAUGEROTTE, *Algebraic elimination of ϵ -transitions*, DMTCS, Volume 7, (2005), pp. 51-70
- [13] DUCHAMP G., FLOURET M., LAUGEROTTE É., LUQUE J-G., *Direct and dual laws for automata with multiplicities* T.C.S. **267**, 105-120 (2001).
- [14] S. EILENBERG, Automata, languages and machines, (Acad. Press, New-York, 1974).
- [15] M. FLIESS, Sur le plongement de l'algèbre des séries rationnelles non commutatives dans un corps gauche. CRAS Ser. A **271** pp 926-927 (1970).
- [16] M. FLIESS, Matrices de Hankel, Jour. of Pure and Appl. Math. **53**, 197-222, (1994).
- [17] I.M. GELFAND, V.S. RETAKH, A theory of non-commutative determinants and characteristic functions of graphs, Funct. Anal. Appl. **26** (1992).
- [18] I.M. GELFAND, D. KROB, A. LASCoux, B. LECLERC B, V.S. RETAKH and J.-Y. THIBON, Noncommutative symmetric functions, Adv. in Math. **112**, 218-348 (1995).
- [19] GOLAN J. S., *Power Algebras over Semirings: With Applications in Mathematics and Computer science*. Kluwer Academic Publishers (1999).

- [20] GOLAN J. S., *Semirings and Affine Equations over Them: Theory and Applications*. Kluwer Academic Publishers (2003).
- [21] A. HEYTING, Die Theorie der linearen Gleichungen in einer Zahlenspezies mit nichtkommutativer Multiplikation, *Math. Ann.*, **98**, 465-490 (1927).
- [22] G. JACOB, Représentations et substitutions matricielles dans la théorie matricielle des semigroupes. Thèse, Univ. de Paris (1975).
- [23] KATRIEL J., DUCHAMP G., *Ordering relations for q-boson operators, continued fractions techniques, and the q-CBH enigma*. *Journal of Physics A* **28** 7209-7225 (1995).
- [24] J. LEWIN, Fields of fractions for group algebras of free groups, *Trans. Amer. Math. Soc.* **192**, 339-346, (1974).
- [25] D.S. PASSMAN, *The algebraic structure of group rings*, John Wiley - Interscience (1977).
- [26] A.R. RICHARDSON Simultaneous linear equations over a division ring. *Proc. Lond. Math. Soc.*, **28**, 395-420 (1928).
- [27] M.P. SCHÜTZENBERGER, On the definition of a family of automata, **4**, *Information and Control*, 275-270, (1961).
- [28] M.P. SCHÜTZENBERGER, On a theorem of R. Jungen, *Proc. Amer. Math. Soc.* **13**, 885-889, (1962).